

Prospects for Non-Fermi-Liquid Behavior of a Two Level Impurity in a Metal.

Aris L. Moustakas and Daniel S. Fisher

Department of Physics, Harvard University, Cambridge, MA 02138

It has been claimed that an impurity hopping between two sites in a metal can lead to non-Fermi liquid behavior if the bias between the two sites is tuned to zero. Here it is shown that several extra tunnelling processes are allowed; those involving bare impurity tunnelling as well as two-electron-assisted tunnelling play important roles. These terms generally drive the system away from the intermediate coupling non-Fermi-liquid fixed point, which has the character of the two channel Kondo problem. In order to find the non-Fermi liquid behavior a combination of these terms must be small over a range of energy scales. The prospects for obtaining these conditions are studied by analyzing both the weak coupling and the intermediate coupling behavior, together with the connecting crossover region. Unfortunately, the conclusions are rather pessimistic. The phase diagram and the important crossover are analyzed via bosonization and refermionization and some of the results compared to those obtained from conformal field theory. Finally, an additional non-Fermi liquid fixed point is found, a multicritical point with two relevant directions even in the unbiased case.

I. INTRODUCTION

A two-channel Kondo impurity in a metal is, perhaps, the most promising impurity system to exhibit low temperature non-Fermi-liquid behavior experimentally. This hope rests on the relatively few degrees of freedom involved (a local spin doublet coupled to two degenerate “flavors” of spin- $\frac{1}{2}$ fermions) with the symmetry group under which they transform being not very complicated [$SU(2)_{\text{flavor}} \times SU(2)_{\text{spin}} \times U(1)_{\text{charge}}$]. As a result one might hope that it would not be difficult to find a system whose low temperature behavior would be described by the two-channel Kondo system.

Yet, for more than a decade since it was first introduced by Nozières and Blandin¹, no experimental realization of this model has been conclusively demonstrated. The difficulty lies in the fact that the non-Fermi-liquid fixed point is unstable to various symmetry-breaking processes which turn out to be present in real experimental situations. For example, Nozières and Blandin¹ pointed out that anisotropy between the two flavor channels, caused by lattice effects, would destroy the non-Fermi-liquid ground state. Since this anisotropy could not be made to vanish for the known cases, the search for non-Fermi-liquid behavior with conventional spin-Kondo systems was suspended. Cox² pointed out, however, that under certain symmetry conditions, local quadrupolar degrees of freedom could result in two channel Kondo-like coupling; unfortunately, dilute impurity systems of this type have proven hard to make.

Later, Vladár and Zawadowski³ suggested that a non-magnetic impurity tunnelling between two sites in a metal could be modeled as a two-channel Kondo system in which the roles of the channels and the spins in the original formulation are interchanged. In this system the spin of the electron plays the role of the “flavor channels”, so that the anisotropy between “channels” is

no longer an issue since in zero external magnetic field the spin-up and the spin-down electrons are degenerate. This led Vladár and Zawadowski³ to predict non-Fermi-liquid behavior in such a system. Recently Ralph *et al.*^{4,5} have interpreted low temperature tunnelling data in very small metallic contacts in terms of two-channel Kondo-like physics. Their measurements are claimed to be consistent with certain exact results obtained by Affleck and Ludwig⁶, but at this point, the interpretation is still controversial⁷.

Unfortunately, the mapping of the two-site impurity to the two-channel Kondo (2CK) system is far from exact, and, even with no anisotropy between “channels” (the spin-up and spin-down electrons) there are other processes present in the tunnelling impurity system which are relevant and hence generically destroy the non-Fermi-liquid ground state. These processes, which cannot be neglected have not been treated adequately in the literature^{3,8,9}.

In this paper we carefully consider the mapping between the tunnelling impurity system and the two-channel Kondo (2CK) problem. We first analyze the behavior when the impurity tunnelling starts to become important as the temperature is lowered; this is analogous to the weak coupling regime of the Kondo problem and is needed in order to understand which tunnelling processes will dominate at low temperatures. We then analyze the intermediate coupling behavior to investigate whether there are parameter regimes in which the system will be governed by the 2CK fixed point over a reasonable range of temperatures – as would be needed to observe the non-Fermi liquid behavior experimentally.

Unfortunately, even in the optimal case of an impurity tunnelling between two identical sites so that the system has an extra Z_2 symmetry, we find that the 2CK fixed point is only accessible if two tunnelling processes, which are very different physically, nearly exactly cancel.

Generically, the system will exhibit Fermi liquid behavior at low temperatures.

We analyze the behavior near the 2CK point in detail, focusing on the connection between the physical operators and those that appear as “natural” operators in the 2CK language. It is found that, on the critical manifold which can be obtained in the symmetric impurity problem by adjusting one parameter, there are *four* leading irrelevant directions in contrast to the behavior for the pure 2CK problem analyzed by Sengupta and Georges¹⁰. Various symmetry breaking terms are also studied and our results recover those derived by Affleck *et al*^{11,12} with conformal field theory techniques.

A somewhat surprising feature emerges in the fuller phase diagram of the symmetric impurity model: a *second* fixed point which exhibits non-Fermi liquid behavior, albeit one with two relevant directions in the Z_2 symmetric case.

A. Outline

In the remainder of the Introduction we motivate the form of the Hamiltonian in which we focus and interpret the various terms that should appear. In Section II we derive an effective Hamiltonian that we will study and analyze its symmetries, while in Section III the weak coupling analysis is outlined. In Section IV the behavior near the intermediate coupling 2CK fixed point and, for completeness, the various symmetry breaking operators near the 2CK fixed point are studied. In Section V we discuss the existence of an extra novel fixed point at intermediate coupling. In Section VI, we discuss the accessibility of the 2CK fixed point and draw our conclusions. Finally, in the Appendices, the details of the weak-coupling (Appendix A) and the intermediate-coupling analysis (Appendix B) are presented; the comparison of our results to those obtained by conformal field theory is made in Appendix C.

B. Physical Picture

The system we wish to describe is an impurity or heavy particle which can hop back and forth between two sites coupled to a bath of electrons. The two sites may or may not be equivalent but we will primarily focus on the symmetric (equivalent) case. The asymmetric case can readily be treated in a similar manner. The effects of the interaction of this impurity with the electrons can be manifested in a number of ways. First, the electrons will tend to screen the charge of the impurity. Thus the impurity will hop between the two sites carrying with it tightly bound electrons which can move fast enough to adjust to the position of the impurity; it is convenient to consider these to simply be part of the “dressed” impurity particle. However, in addition, as the impurity

moves it may also redistribute the low energy electronic excitations near the Fermi surface. Since we are interested in the low energy physics, these processes must be treated directly. For simplicity we consider only *s*-wave scattering off the impurity.

If an impurity of charge Ze hops between two well separated sites, then the Friedel sum rule relates the (*s*-wave) scattering phase shift off a static impurity, δ , to the electronic charge that will be moved to screen the impurity as it moves adiabatically from one site to the other, via $Z = 2\delta/\pi$, with the factor of two due to the two spin species. Conversely, if the two sites are close together, one can still usefully speak of an effective charge Q (per spin) which plays an analogous role to $Q = \delta/\pi$ in the well separated case, but is no longer simply related either to scattering phase shifts or to the impurity charge. It will instead turn out to be exactly the “orthogonality catastrophe” exponent that determines the system size dependence of the overlap between the electronic ground states with the impurity at the two sites.¹³ For simplicity, we will focus on Q in the range $0 \leq Q \leq 1$, corresponding to repulsive interactions. (As shown in reference 14, other ranges of the effective charge can be reduced to this case via a set of more complicated combined impurity-electron processes related to those we consider here.)

The important processes, in addition to hopping of the (dressed) impurity by itself, will be those in which one or two low energy electrons move in the *opposite* direction to that the impurity hops. These processes can, for $Q > 1/2$, reduce the effective charge that must relax to the new impurity position, thereby decreasing the orthogonality between the pre- and post-hop configuration; this results in a larger amplitude for the combined impurity-electron process at low temperatures, relative to the simple impurity hop process. [Note that in reference 14, a sign error in the definition of Q resulted in the incorrect interpretation being given to these processes; this error does not affect the conclusions, just the interpretation of the combined hopping processes].

In order to proceed with the analysis of the low temperature behavior of interest it would appear to be important to assess the relative magnitudes of the amplitudes, at some high energy scale, of the processes discussed above. However, it has been claimed¹⁷ that this is essentially impossible for strong electron-impurity interactions. Indeed Kagan and Prokof'ev¹⁷ have claimed that a sensible Hamiltonian cannot be written in terms of a simple two level system since the high energy electronic degrees of freedom cannot be properly taken into account. Although there are indeed real difficulties here, it should nevertheless be possible to introduce *effective* amplitudes at some intermediate energy scale and then analyze the behavior of the system in the phase space of these effective parameters. In general, unless there are specific reasons to prevent it, one would expect that most combinations of parameters could, in principle, be realized. We thus approach the problem via this route

and start with an effective Hamiltonian at an intermediate energy scale, at, say, some fraction of the conduction bandwidth. As we shall see, the extra hopping processes will in any case be generated at low energies.

II. EFFECTIVE HAMILTONIAN

The important electronic degrees of freedom at low energies are those that interact with the impurity in one of its two positions “1” and “2”. These are just the s -wave conduction electrons around the positions of the two sites. Thus at each energy, there will be two important electronic degrees of freedom per spin. However, unless the two sites are very far apart (in which case the impurity tunnelling rates will be negligible and hence not of interest), the two sets of s -wave electrons will *not* be orthogonal; this will play an important role in what follows. If we label the two sets of s -wave electrons by their energy, ϵ , measured from the Fermi surface, then for each ϵ there is an (essentially) unique pair of linear combinations of the two s -wave states, that are orthonormal and transform into each other under interchange of the two sites. We label the annihilation operators of this orthonormal pair $c_{1\epsilon}$ and $c_{2\epsilon}$ with anticommutation relations

$$\{c_{i\epsilon}, c_{j\epsilon'}^\dagger\} = 2\pi\delta(\epsilon - \epsilon')\delta_{ij} \quad (1)$$

with the “1” and “2” denoting the sites near which the wavefunction is larger. The impurity interacts with, simply, the operators

$$c_{1,2} = \int \frac{d\epsilon}{2\pi} c_{1,2\epsilon}, \quad (2)$$

although in each position the impurity will couple to *both* c_1 and c_2 due to the non-orthogonality of the two electronic s -wave states. With time reversal invariance, which we assume henceforth, the most general interaction with the impurity becomes (ignoring spin for now)

$$\begin{aligned} U = d_1^\dagger d_1 & \left[V_1 (c_1^\dagger c_1 + c_2^\dagger c_2) + V_2 (c_1^\dagger c_2 + c_2^\dagger c_1) \right. \\ & \left. + V_3 (c_1^\dagger c_1 - c_2^\dagger c_2) \right] \\ + d_2^\dagger d_2 & \left[V_1 (c_2^\dagger c_2 + c_1^\dagger c_1) + V_2 (c_2^\dagger c_1 + c_1^\dagger c_2) \right. \\ & \left. + V_3 (c_2^\dagger c_2 - c_1^\dagger c_1) \right] \end{aligned} \quad (3)$$

where $d_{1,2}$ are the annihilation operator of the impurity at the sites. Using the obvious identity $d_1^\dagger d_1 + d_2^\dagger d_2 = 1$, Eq(3) can be written as

$$\begin{aligned} U = V_1 & (c_1^\dagger c_1 + c_2^\dagger c_2) + V_2 (c_1^\dagger c_2 + c_2^\dagger c_1) \\ + V_3 & (d_1^\dagger d_1 - d_2^\dagger d_2) (c_1^\dagger c_1 - c_2^\dagger c_2). \end{aligned} \quad (4)$$

Note the appearance of an effective electronic hopping term V_2 , caused by the scattering by the impurity; this will vanish if the sites are far apart, but in general will be comparable to the other terms.

The first term in Eq(4) which is the average of the interaction over the two impurity positions, merely produces a constant phase shift for both “1” and “2” electrons at the Fermi level and, combined with other operators, gives rise only to irrelevant terms. Therefore we will ignore it at this point although in Section IV an irrelevant operator it gives rise to will play a role in our discussion of the intermediate coupling behavior.

With the effective hopping charge Q in the range $[0, 1]$, there are three hopping processes that must be considered¹⁴:

$$\begin{aligned} \mathcal{H}_{hop} = d_2^\dagger d_1 & \left[\Delta_0 + \frac{\Delta_1}{2} (c_{1\uparrow}^\dagger c_{2\uparrow} + c_{1\downarrow}^\dagger c_{2\downarrow}) \right. \\ & \left. + \Delta_2 c_{1\uparrow}^\dagger c_{2\uparrow} c_{1\downarrow}^\dagger c_{2\downarrow} \right] + h.c. \end{aligned} \quad (5)$$

representing hopping of the (dressed) impurity, jointly with, respectively, 0, 1 and 2 electrons moving the opposite way. Although we might start at a high energy scale with negligible Δ_1 and Δ_2 , these will be generated under renormalization and hence must be included.

In order to analyze the renormalization group flows, it is convenient to approximate the conduction band s -wave electrons $c_{1\epsilon}$ and $c_{2\epsilon}$ by a linear dispersion with a cutoff, at short times, τ_c , roughly the inverse bandwidth. Then the interactions with the impurity will essentially be replaced by corresponding phase shifts, specifically V_3 replaced by an effective phase shift that we denote πQ_0 ; Q_0 will have the interpretation of an effective “charge”. We then have, after inserting powers of τ_c to make the couplings dimensionless and factors of π for convenience,

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_{int} + \mathcal{H}_{hop} \quad (6)$$

with

$$\mathcal{H}_0 = \sum_\sigma \int \frac{d\epsilon}{2\pi} \epsilon \left[c_{1\sigma\epsilon}^\dagger c_{1\sigma\epsilon} + c_{2\sigma\epsilon}^\dagger c_{2\sigma\epsilon} \right], \quad (7)$$

$$\begin{aligned} \mathcal{H}_{int} = \pi Q_0 & (d_2^\dagger d_2 - d_1^\dagger d_1) \sum_\sigma (c_{2\sigma}^\dagger c_{2\sigma} - c_{1\sigma}^\dagger c_{1\sigma}) \\ + \pi y & \sum_\sigma (c_{1\sigma}^\dagger c_{2\sigma} + c_{2\sigma}^\dagger c_{1\sigma}) \end{aligned} \quad (8)$$

and

$$\begin{aligned} \mathcal{H}_{hop} = d_2^\dagger d_1 & \left[\frac{\Delta_0}{2\pi\tau_c} + \frac{\Delta_1}{2} \sum_\sigma c_{1\sigma}^\dagger c_{2\sigma} \right. \\ & \left. + \Delta_2 2\pi\tau_c c_{1\uparrow}^\dagger c_{2\uparrow} c_{1\downarrow}^\dagger c_{2\downarrow} \right] + h.c. \end{aligned} \quad (9)$$

We have rescaled the electronic term V_2 to a coefficient y which will play the role of a “fugacity” for electronic hops.

A. Symmetries

It is important at this stage to examine the symmetries of Eq(6). In addition to time reversal, conservation of the electrons ($c \rightarrow e^{i\phi}c$), conservation of the impurity ($d \rightarrow e^{i\phi}d$), and $SU(2)$ spin symmetry, the only other symmetry is interchange of the two sites and the corresponding electronic states ($1 \leftrightarrow 2$). Note, however, that if the only hopping term had been Δ_1 , and if y vanished, there would be an *extra* artificial symmetry $d_1 \rightarrow e^{i\phi}d_1$, $c_1 \rightarrow e^{i\phi}c_1$, $d_2 \rightarrow d_2$, $c_2 \rightarrow c_2$ corresponding to conservation of $N_1 = d_1^\dagger d_1 + n_{c_1}$ and similarly N_2 separately, where n_{c_1} is the number of the “one” electrons which, in the absence of the channel mixing term y , are independent of the “two” electrons. As shown in reference 14, even if the electronic states had been optimally chosen so that there was no mixing of “one” and “two” electrons at the Fermi energy, the energy dependence of scattering off the impurity would generate extra mixing terms in $\mathcal{H}_{int}$, that cannot simply be expressed in terms of c_1 and c_2 . These would break the artificial symmetry and under renormalization generate a $y (c_1^\dagger c_2 + h.c.)$ mixing term even in the absence of impurity motion. Thus it is best to include y from the beginning. (The neglected energy dependent scattering terms will then not play an important role). In order to understand the difficulties of reaching the 2CK-like fixed point, this step is *crucial*.

The artificial symmetry in the absence of the channel mixing and Δ_0 , Δ_2 terms, corresponds to a conserved pseudo-spin $N_1 - N_2$ which is the sum of the “ z -components” of the impurity pseudo-spin $d_2^\dagger d_2 - d_1^\dagger d_1$ and an electronic pseudo-spin $n_{c_2} - n_{c_1}$. This pseudo-spin can play the role of spin for the two-channel Kondo effect and, indeed, under renormalization the system will flow to this intermediate coupling 2CK fixed point if $Q > 0$.¹⁵ Unfortunately, there is no natural small parameter which keeps the pseudo-spin symmetry breaking terms small.¹⁶

B. Bosonization

In order to carry out the renormalization group analysis for small bare impurity hopping rates, it is useful, as is standard, to bosonize the electronic degrees of freedom, treating the electronic states $c_{1,2\epsilon}$ as those of a one-dimensional system with two sets of right moving electrons with “wavevectors” $v_F(k - k_F) \propto \epsilon$. It is simplest to set the Fermi velocity, $v_F = 1$, and treat ϵ like a wavevector index, defining

$$c_{j\sigma}(x) \equiv \int \frac{d\epsilon}{2\pi} e^{i\epsilon x} c_{j\sigma\epsilon} = \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_{j\sigma}(x)} \quad (10)$$

so that

$$c_{j\sigma}^\dagger(x) c_{j\sigma}(x) = \frac{1}{2\pi} \frac{\partial \Phi_{j\sigma}(x)}{\partial x} \quad (11)$$

with $j = 1, 2$ $\sigma = \uparrow, \downarrow$ and $\Phi_{j\sigma}$ being the corresponding bosonic degrees of freedom, where we have followed Emery and Kivelson’s notation.¹⁸ Only $c_{j\sigma} \equiv c_{j\sigma}(x=0)$ couples to the impurity. Note that in the standard expression Eq(11) the left hand side is normal ordered and therefore the (infinite) uniform charge density does not appear. Also corrections that vanish as $\tau_c \rightarrow 0$ are neglected; we will be careful to include the effects of extra terms when they play an important role.

Since we will later need to be careful to have the proper anticommutation relations, we must insert extra factors of the form $\exp(i\pi N_\mu)$, with $N_\mu \equiv \int dx \Psi_\mu^\dagger(x) \Psi_\mu(x)$, into some of the bosonized expressions to ensure anticommutations of the different Fermi fields. These will not play a role as long as no spin-flip processes occur, and, for the time being we ignore them; the needed modifications are spelled out in Appendix B.

It is useful to define even and odd components of the Bose fields

$$\Phi_{e,o\sigma} = \frac{1}{\sqrt{2}} (\Phi_{2\sigma} \pm \Phi_{1\sigma}), \quad (12)$$

the Hamiltonian then becomes

$$\begin{aligned} \mathcal{H} = \mathcal{H}_0 &+ \frac{Q_0}{\sqrt{2}} \sigma_z \sum_\sigma \frac{\partial \Phi_{o\sigma}}{\partial x} + y \sum_\sigma \cos \Phi_{o\sigma} \\ &+ \frac{\Delta_0}{2\pi\tau_c} \sigma_x + \frac{\Delta_1}{4\pi\tau_c} \sum_\sigma \left(\sigma_+ \exp \left[i\sqrt{2}\Phi_{o\sigma} \right] + h.c. \right) \\ &+ \frac{\Delta_2}{2\pi\tau_c} \left(\sigma_+ \exp \left[i\sqrt{2}(\Phi_{o\uparrow} + \Phi_{o\downarrow}) \right] + h.c. \right) \end{aligned} \quad (13)$$

where in all the coupling terms the bosonic fields are evaluated at $x = 0$ and we use the impurity pseudo-spin operators

$$\sigma_z = d_2^\dagger d_2 - d_1^\dagger d_1 \quad (14)$$

and

$$\sigma_+ = d_2^\dagger d_1, \quad \sigma_- = d_1^\dagger d_2. \quad (15)$$

The conduction electron part of the Hamiltonian can be written in terms of boson creation and annihilation operators $\phi_\mu(\epsilon)$ with canonical commutation relation

$$[\phi_\mu(\epsilon), \phi_\nu^\dagger(\epsilon')] = 2\pi \delta_{\mu\nu} \delta(\epsilon - \epsilon') \quad (16)$$

via

$$\Phi_\mu(x) = \int_0^\infty \frac{d\epsilon}{\sqrt{2\pi\epsilon}} [\phi_\mu(\epsilon) e^{i\epsilon x} + \phi_\mu^\dagger(\epsilon) e^{-i\epsilon x}] e^{-\frac{\epsilon\tau_c}{2}} \quad (17)$$

and

$$\mathcal{H}_0 = \sum_\mu \int_0^\infty \frac{d\epsilon}{2\pi} \epsilon \phi_\mu^\dagger(\epsilon) \phi_\mu(\epsilon) e^{-\epsilon\tau_c} \quad (18)$$

which involves positive energy parts only. Here τ_c^{-1} is the energy cutoff and μ represents the various Bose fields, i.e. for Eq(13), $\mu = (e \uparrow, e \downarrow, o \uparrow, o \downarrow)$. We see that Eq(13) does not include any terms with $\Phi_{e\downarrow}$ or $\Phi_{e\uparrow}$. Thus, up to operators that are irrelevant for weak coupling, the impurity is decoupled from the even boson fields.

It is convenient, following Emery and Kivelson¹⁸, to decompose the odd field Φ_o which couples to the impurity, into a spin and charge part, by

$$\Phi_{o\sigma} = \frac{1}{\sqrt{2}} (\Phi_c + \sigma \Phi_s) \quad (19)$$

with $\sigma = \pm$ for spin $\uparrow, \downarrow$, respectively. The second term in Eq(13) becomes simply $Q_0 \sigma_z \frac{\partial \Phi_c}{\partial x} (0)$. This term, which represents the difference in phase shifts for electron scattering off the two positions of the impurity, can be shifted away by a unitary transformation which changes the naive weak coupling scaling of the hopping terms; this is the conventional approach used^{9,10,14,18} to derive the weak coupling flows discussed in the next section.

Although the even fields will not play much role for the time being, for later purposes we also introduce even fields Φ_{ec}, Φ_{es} by

$$\Phi_{e\sigma} = \frac{1}{\sqrt{2}} (\Phi_{ec} + \sigma \Phi_{es}). \quad (20)$$

Note that any sum or difference of any two of the $\Phi_{j\sigma}$, i.e. those that appear from operators bilinear in electron operators, can be written as a sum or difference of two of the fields $\Phi_s, \Phi_c, \Phi_{es}, \Phi_{ec}$ with coefficients of *unity*; this enables the method of Emery and Kivelson¹⁸ to work.

III. WEAK HOPPING ANALYSIS

In order to connect the various amplitudes at the relatively high energy scale of the effective Hamiltonian (Eq(13)) to their renormalized values at low energies we must analyze the weak coupling renormalization group (RG) flow equations for the amplitudes in Eq(13). The magnitudes of the various terms at the crossover scale to intermediate coupling will determine which regions of the initial parameter space can flow near to the 2CK fixed point.

Following the procedure in reference 14 we transform $\mathcal{H}$ to $U\mathcal{H}U^\dagger$ using the unitary operator

$$U = \exp[-i\sigma_z Q_0 \Phi_c] \quad (21)$$

Subsequently we follow the RG approach described there and obtain the following flow equations for the various amplitudes, where for later convenience we introduce

$$q = \frac{1}{2} - Q_0, \quad (22)$$

which lies in the range $(-1/2, 1/2)$:

$$\begin{aligned} \frac{d\Delta_0}{dl} &= \left(\frac{1}{2} + 2q - 2q^2\right) \Delta_0 + y\Delta_1 + O(\Delta^3) \\ \frac{d\Delta_2}{dl} &= \left(\frac{1}{2} - 2q - 2q^2\right) \Delta_2 + y\Delta_1 + O(\Delta^3) \\ \frac{d\Delta_1}{dl} &= \left(\frac{1}{2} - 2q^2\right) \Delta_1 + 2y(\Delta_0 + \Delta_2) + O(\Delta^3) \\ \frac{dq}{dl} &= -2q \left(\Delta_0^2 + \Delta_2^2 - \frac{1}{2}\Delta_1^2\right) + (\Delta_0^2 - \Delta_2^2) \\ \frac{dy}{dl} &= \Delta_1(\Delta_0 + \Delta_2) \end{aligned} \quad (23)$$

The important cross-terms in the first three equations in Eq(23) that are proportional to the electronic mixing term y , have a simple physical interpretation: they represent the effects of an impurity and an electronic hop both occurring within a short time interval so that, at lower energies, this appears as simply the corresponding combined process. Note that we have not included $O(y^2)$ terms in the above equations; the definition of these will depend on the RG procedure, and they will not qualitatively change the behavior. Thus, in the spirit of focusing on the important processes and terms, we ignore them¹⁹.

Noting that q and y are constant to order $O(\Delta^2)$, to analyze the flow for weak hopping, we can safely set them to their initial values, q_0 and y_0 . Then the first three equations can be diagonalized exactly; the details are discussed in Appendix A. The RG eigenvalues for the hopping terms, about the zero hopping fixed line are

$$\begin{aligned} \lambda_{\pm} &= \frac{1}{2} - 2q_0^2 \pm 2\sqrt{q_0^2 + y_0^2} \\ \lambda_0 &= \frac{1}{2} - 2q_0^2. \end{aligned} \quad (24)$$

We now note that for $0 \leq Q_0 \leq 1$, corresponding to $|q_0| \leq \frac{1}{2}$, at least two eigenvalues are positive so that impurity hopping is always relevant (leading to the conclusion of *absence* of impurity localization¹⁴); likewise for other ranges of Q_0 , there will always be at least two relevant hopping processes if there is only s -wave scattering off the impurity.

The Kondo temperature, T_K , is the energy scale at which the first of the impurity hopping processes becomes of order unity—i.e. of order the renormalized bandwidth.

The system considered by Vladár and Zawadowski³ and Vladár *et al*⁸, essentially amounts to neglecting Δ_2 and y ; for small $|q|$, which will turn out to be the most interesting case, this misses part of the physics. The reason is simply that they neglect one relevant operator Δ_2 which mixes with the other two to give the correct eigenvalues (Eq24). Furthermore, as will be seen later, non-vanishing values of Δ_2 and y are crucial to give the correct renormalization flows close to the intermediate coupling fixed point.

If we only kept Q and Δ_1 non-zero, their weak coupling flows would be (up to coefficients) like those for J_z

and $J_\perp$ for the conventional Kondo problem. The Kondo scale is then simply

$$\frac{T_K}{W} \sim \left(\frac{\Delta_1}{W} \right)^{\frac{1}{\lambda_0}}, \quad (25)$$

after reinserting factors of the bandwidth, W . (For the special value $\Delta_1 = 2Q$, $T_K \sim e^{-\frac{1}{\Delta_1}}$ like in the well known anti-ferromagnetic Heisenberg Kondo problem). For this artificial case, at scales below T_K the novel two channel Kondo physics will indeed appear, as we will discuss later. This results from the approach, at low energies, of the system to the intermediate coupling 2CK fixed point.

Unfortunately, the breaking of the artificial pseudo-spin symmetry leads to the appearance of terms which generically drive the flow *away* from the 2CK fixed point. In order to analyze whether the system can get near to the 2CK fixed point — the prerequisite for observation of non-Fermi liquid behavior —, we must be able to identify the operators near the 2CK fixed point in terms of the original terms in the Hamiltonian; the magnitude of the operators, in particular the relevant ones, can then be determined, roughly, by “matching” the coefficients at the crossover scale, T_K , between the weak and intermediate coupling regimes.

From Appendix A, we see that the crossover temperature, T_K , will generally be a complicated function of the original parameters. Before examining the magnitude of the various important terms at T_K , we turn to the behavior near the 2CK fixed point; this will tell us which terms need to be small at scales of order T_K for 2CK behavior to obtain.

IV. INTERMEDIATE COUPLING ANALYSIS AND TWO CHANNEL KONDO FIXED POINT

From the weak coupling flow equations in Eq(23), it is apparent that, in the absence of electronic mixing ($y=0$) there is a special value of the effective impurity charge Q : for $Q = \frac{1}{2}$, corresponding to $q = 0$, Δ_0 , Δ_1 and Δ_2 all scale in the same way with eigenvalue $\lambda = \lambda_0 = \lambda_\pm = \frac{1}{2}$. By analogy to the Toulouse limit of the conventional one channel Kondo problem, it is thus natural to look for a solvable point that corresponds to $y = q = 0$, inspired by the observation that free Fermi fields have scaling dimension of $\frac{1}{2}$ and thus might be used to represent all of the hopping terms that appear for $q = 0$. This has been carried out recently by Emery and Kivelson¹⁸ who “refermionize” the bosonized operators that appear in the Hamiltonian enabling the computation of the scaling of the various important operators.

By examining the weak coupling flows, it is apparent that special behavior might occur when $\Delta_0 + \Delta_2 = 0$. Defining

$$\Delta_+ \equiv \Delta_0 + \Delta_2, \quad (26)$$

we see that, at least to the order included in Eq(23), for $y = 0$ and $\Delta_+ = 0$, q flows to zero, y is *not* generated and one might hope that the flow would be towards the 2CK fixed point. Indeed, the intermediate coupling analysis shows that this can occur, even if

$$\Delta_- \equiv \Delta_0 - \Delta_2 \quad (27)$$

is non-zero so that the artificial pseudo-spin symmetry is broken.

Physically the role of Δ_+ is very surprising. The processes represented by Δ_0 and Δ_2 are very different and the definitions which make them dimensionless are clearly cutoff dependent. Thus the special critical behavior must not in general occur exactly at $\Delta_+ = 0$, since the location — but not the existence — of the critical manifold will be affected by irrelevant operators. In particular, from the weak coupling flows we can see that a non-zero q combined with $\Delta_- \neq 0$ *will* generate Δ_+ , thus a “bare” Δ_+ that is non-zero will be needed for the flow to go to the 2CK fixed point asymptotically.

In order to understand the intermediate coupling behavior and the special role of $Q = \frac{1}{2}$, following Emery and Kivelson¹⁸ and the analogous Toulouse limit²⁰ of the conventional Kondo problem, we perform a unitary transformation with

$$U = \exp \left[-\frac{i}{2} \sigma_z \Phi_c \right] \quad (28)$$

which transforms the Hamiltonian to $\tilde{\mathcal{H}} = U\mathcal{H}U^\dagger$

$$\begin{aligned} \tilde{\mathcal{H}} = & \mathcal{H}_0 + \frac{\Delta_1}{2\pi\tau_c} \sigma_x \cos \Phi_s + \frac{\Delta_-}{2\pi\tau_c} \sigma_y \sin \Phi_c \\ & + \frac{\Delta_+}{2\pi\tau_c} \sigma_x \cos \Phi_c + 2y \cos \Phi_c \cos \Phi_s - q\sigma_z \frac{\partial \Phi_c}{\partial x} \\ & + \frac{u}{\pi} \frac{\partial \Phi_{ec}}{\partial x} - \frac{w}{2\pi} \sigma_x \cos \Phi_c \frac{\partial \Phi_{ec}}{\partial x} \end{aligned} \quad (29)$$

where we have combined the terms

$$\begin{aligned} & \frac{\Delta_2}{2\pi\tau_c} (\sigma_+ e^{i\Phi_c} + \sigma_- e^{-i\Phi_c}) + \frac{\Delta_0}{2\pi\tau_c} (\sigma_+ e^{-i\Phi_c} + \sigma_- e^{i\Phi_c}) \\ & = \frac{\Delta_-}{2\pi\tau_c} \sigma_y \sin \Phi_c + \frac{\Delta_+}{2\pi\tau_c} \sigma_x \cos \Phi_c, \end{aligned} \quad (30)$$

and abbreviated $\Phi_\mu (x=0)$ simply by Φ_μ . We have also reintroduced some of the coupling terms to the even fields, in particular the marginal term $u \frac{\partial \Phi_{ec}}{\partial x}$ that arises from the impurity-position independent part of the electron-impurity scattering (V_1 in Eq(4)) and the term $w \sigma_x \cos \Phi_c \frac{\partial \Phi_{ec}}{\partial x}$, which arises from the combination of u and impurity hopping terms Δ_+ and is irrelevant for weak hopping, will play roles in our analysis. An additional irrelevant term, $\sigma_y \sin \Phi_s \frac{\partial \Phi_{ec}}{\partial x}$ couples the impurity to the electronic *spin* degrees of freedom, but does not feed back to the other operators and thus we ignore it for the time being. Its effect will be discussed further in Appendix C.

A. Symmetries

Since we expect that the symmetries will play an important role, we should examine what the original symmetries correspond to in $\tilde{\mathcal{H}}$ and ensure that there are no extra symmetries which might have arisen from the discarding of operators which were naively irrelevant, since, as shown in reference 14, such procedures, especially when combined with “large” transformations, such as Eq(28), can be dangerous.

1. Gauge invariance and spin conservation

Since there are no spin-flip processes, separate gauge transformations can be made for each spin $\Phi_{j\sigma}(x) \rightarrow \Phi_{j\sigma}(x) + \theta_\sigma$, corresponding to $\Phi_{e\sigma} \rightarrow \Phi_{e\sigma} + \sqrt{2}\theta_\sigma$. This does not play much role as the Φ_{ec} -field enters only as a derivative $\frac{\partial \Phi_{ec}}{\partial x}$ and the Φ_{es} -field decouples from the impurity at the level at which we work (there is feedback, under renormalization from other operators involving Φ_e but these only modify pre-existing terms in $\tilde{\mathcal{H}}$). However, if the z -component of electron spin is *not* conserved, then only the symmetry $\Phi_{ec} \rightarrow \Phi_{ec} + \theta_{ec}$ remains.

2. Periodicity

The definition of the Fermi fields, Eq(10), implies that shifting any $\Phi_{j\sigma}$ by 2π should leave $\tilde{\mathcal{H}}$ unchanged. Depending on whether one or both spin components are shifted, this implies that (ignoring shifts in Φ_{ec})

$$\Phi_c \rightarrow \Phi_c + 2\pi \quad (31)$$

and

$$\Phi_s \rightarrow \Phi_s + 2\pi \quad (32)$$

are independent symmetries, as is the combination

$$\begin{aligned} \Phi_c &\rightarrow \Phi_c + \pi, \quad \Phi_s \rightarrow \Phi_s + \pi \\ \sigma_x &\rightarrow -\sigma_x, \quad \sigma_y \rightarrow -\sigma_y; \end{aligned} \quad (33)$$

with the necessity for the simultaneous transformation of $\sigma_{x,y}$ resulting from the unitary transformation, U , which involves Φ_c so that $\Phi_c \rightarrow \Phi_c + \pi$ introduces an extra $\exp(-\frac{\pi i}{2}\sigma_z) = -i\sigma_z$ factor into U yielding $\sigma_{x,y} \rightarrow -\sigma_{x,y}$ in $\tilde{\mathcal{H}}$.

3. Interchange

Interchanging sites one and two is equivalent to

$$\begin{aligned} \Phi_c &\rightarrow -\Phi_c, \quad \Phi_s \rightarrow -\Phi_s, \\ \sigma_y &\rightarrow -\sigma_y, \quad \sigma_z \rightarrow -\sigma_z \end{aligned} \quad (34)$$

with $\Phi_{ec,es}$ unchanged.

4. Spin reversal

Flipping electron spins is simply $\Phi_s \rightarrow -\Phi_s$ and $\Phi_{es} \rightarrow -\Phi_{es}$.

5. Time reversal

Time reversal transformations change ingoing to outgoing waves, thereby yielding $x \rightarrow -x$, $i \rightarrow -i$, all $\Phi_\mu(x) \rightarrow -\Phi_\mu(-x)$ and $\sigma_y \rightarrow -\sigma_y$. Note that here we are *not* time reversing the spins.

6. Artificial extra symmetries

We now see that, indeed, $\tilde{\mathcal{H}}$ in Eq(29) with all coefficients non-zero, does *not* have any artificial symmetries. But as seen earlier, an artificial extra pseudo-spin symmetry is possible: pseudo-spin conservation mod 2 corresponds to $\Phi_c \rightarrow \Phi_c + \pi$, and more generally full pseudo-spin symmetry corresponds to independence of the “one” and “two” electrons, i.e. $\Phi_c \rightarrow \Phi_c + \theta_c$ with any θ_c . The terms w , Δ_+ , Δ_- and y all violate this, and it can readily be seen in the representation of $\tilde{\mathcal{H}}$ of Eq(29) that y combined with Δ_- generates Δ_+ , and q combined with Δ_- generates Δ_+ , as expected.

In the representation of Eq(29) we see that there is another possible artificial symmetry: If w , Δ_+ , y and q are all zero, then

$$\Phi_c \rightarrow \pi - \Phi_c \quad (35)$$

becomes a symmetry. This, as we shall see, restricts the system automatically to the stable critical manifold of the 2CK fixed point. But note that because of the unitary transformation of Eq(28) $\Phi_c \rightarrow \pi - \Phi_c$ does *not* correspond to a realizable symmetry in terms of the original variables since it mixes hops involving the impurity alone and those involving the impurity together with two electrons. (Indeed many other irrelevant terms neglected in $\tilde{\mathcal{H}}$ will also violate this artificial symmetry.)

B. Reformionization

If Δ_+ , q , y and the other operators neglected in $\tilde{\mathcal{H}}$ all vanish, then the system will still exhibit the novel intermediate coupling 2CK behavior. As we shall see, the only relevant operator near the 2CK fixed point, which is consistent with the true symmetries of the impurity hopping between two equivalent sites, is Δ_+ , thus only one combination of physical quantities needs to be adjusted to obtain the 2CK behavior. Unfortunately, this is a combination which is not naturally small.

The behavior near the 2CK fixed point can most easily be found following Emery and Kivelson¹⁸ by “reformionizing” the Bose fields Φ_c , Φ_s and Φ_{ec} that appear in $\mathcal{H}$ in Eq(29), noting that $\exp i\Phi_\mu$, (with Φ_μ properly normalized) is like some pseudo-Fermi field. The details are discussed in Appendix B.

Crudely, for $\mu = c, s, ec, es$, each field $e^{i\Phi_\mu}$ is replaced by a new Fermi field, Ψ_μ , and σ_- by a local Fermi field d , with appropriate factors of $e^{i\pi N_\mu}$ and $e^{i\pi N_d}$ to give the correct anticommutation relations. The symmetries can most easily be seen, and the Hamiltonian simplified, by writing the new Fermi fields in terms of a set of Majorana (hermitian) fermions:

$$d = \frac{1}{\sqrt{2}}(\gamma + i\delta) \quad (36)$$

$$\Psi_\mu(x=0) = \frac{1}{\sqrt{2}}(\alpha_\mu + i\beta_\mu).$$

Note that

$$\sigma_z = 2i\gamma\delta. \quad (37)$$

The symmetry restrictions can now be examined in terms of these variables; the details are given in Appendix B. The periodicity of the Bose fields Φ_c and Φ_s simply implies that only terms with an even number of Fermi fields can appear in the Hamiltonian. *Gauge invariance* implies that Ψ_{ec} can only appear as $\Psi_{ec}^\dagger \Psi_{ec}$ and the *z-component of spin conservation* that Ψ_{es} cannot appear in the absence of magnetic fields in the *x*- or *y*-direction. *Spin reversal*, because of the role of the ordering operators, takes $\Psi_s \rightarrow -\Psi_s^\dagger$ and $\Psi_{es} \rightarrow -\Psi_{es}^\dagger$, implying that α_s and α_{es} by themselves are excluded by spin reversal symmetry. *Interchange symmetry* takes $\Psi_c \rightarrow -\Psi_c^\dagger$, $\Psi_s \rightarrow -\Psi_s^\dagger$ and $d \rightarrow d^\dagger$ thereby requiring that α_c and δ must appear together. Finally, *time reversal* takes $x \rightarrow -x$, $i \rightarrow -i$ and $\Phi \rightarrow -\Phi$, allowing only real coefficients of Ψ_μ operators, and hence forbidding terms like $i\gamma\alpha$. The Hamiltonian becomes

$$\begin{aligned} \hat{\mathcal{H}} = \mathcal{H}_0 &+ \frac{i}{\sqrt{2\pi\tau_c}} (\Delta_1\gamma\beta_s + \Delta_- \gamma\beta_c + \Delta_+ \delta\alpha_c) \quad (38) \\ &- 4\pi y\gamma\delta\alpha_c\beta_s + 4\pi q\gamma\delta\alpha_c\beta_c \\ &+ 2iu\alpha_{ec}\beta_{ec} + w\sqrt{2\pi\tau_c}\delta\alpha_c\alpha_{ec}\beta_{ec} \end{aligned}$$

with $\mathcal{H}_0$ the kinetic energy of the four (eight Majorana) new Fermi fields, Ψ_μ . With the full symmetries of the

system, the other five fields (α_s and the *ec*-, *es*-fields), cannot appear in the couplings to the impurity, except in relatively innocuous forms, involving the simple potential coupling to the average position of the impurity, $i\alpha_{ec}\beta_{ec}$ and combinations of this with other terms, as well as the irrelevant term $\delta\alpha_s\alpha_{es}\beta_{es}$, which will be discussed in Appendix C. Note that other potentially important operators, like $\delta\beta_s\alpha_c\beta_c$ and $i\delta\frac{\partial\alpha_c(0)}{\partial x}$ are excluded by time reversal invariance.

The original 2CK problem studied by Emery and Kivelson¹⁸ and Sengupta and Georges¹⁰ corresponds to $\Delta_- = \Delta_+ = y = w = 0$. In our case non-zero Δ_- can be important by observing that “half” of the impurity, γ , couples to both β_c and β_s , thus it is convenient to rediagonalize and make linear combinations of these, β_I and β_X yielding, with

$$\Delta_K = \sqrt{\Delta_1^2 + \Delta_-^2}, \quad (39)$$

$$\begin{aligned} \hat{\mathcal{H}} = \mathcal{H}_0 &+ \frac{i}{\sqrt{2\pi\tau_c}} (\Delta_K\gamma\beta_I + \Delta_+\delta\alpha_c) \quad (40) \\ &+ 4\pi\bar{q}\gamma\delta\alpha_c\beta_X + 4\pi\bar{y}\gamma\delta\alpha_c\beta_I \\ &+ 2iu\alpha_{ec}\beta_{ec} + \sqrt{2\pi\tau_c}w\delta\alpha_c\alpha_{ec}\beta_{ec}, \end{aligned}$$

where $\bar{y}$ and $\bar{q}$ are linear combinations of the original y and q (see Eq(B7) in Appendix B for details). The above rediagonalization of β_c and β_s roughly corresponds to a rotation of “spin” axes in the conventional 2CK language.

From the electronic kinetic energy $\mathcal{H}_0$, the α ’s and β ’s all scale, with time scale τ , as $\tau^{-\frac{1}{2}}$. If all the couplings are small, then the anti-commutation relations of γ and δ imply that they are dimensionless so that Δ_+ and Δ_K scale as $\tau^{-\frac{1}{2}}$, while $\bar{q}$ and $\bar{y}$ are marginal as from the weak coupling analysis of Section II (Eq(23)) and w is irrelevant.

C. Two channel Kondo fixed point and flows

When $\bar{y} = \bar{q} = \Delta_+ = w = 0$, the Hamiltonian in Eq(40) corresponds to the Toulouse limit analyzed by Emery and Kivelson¹⁸. As a free fermion system it can be analyzed straightforwardly. In this limit “half” of the impurity, δ , is uncoupled from the electrons and thus has no dynamics, while the other “half”, γ , gets dynamics from coupling to the electrons with correlations at large imaginary times

$$\langle T_\tau \gamma(\tau) \gamma(0) \rangle \sim \frac{1}{\tau}. \quad (41)$$

Together these yield the non-Fermi liquid 2CK behavior

$$\langle T_\tau \sigma_z(\tau) \sigma_z(0) \rangle \sim \frac{1}{\tau}. \quad (42)$$

It is important to note that²¹ the solvable Hamiltonian is *not* generally at the 2CK fixed point. Indeed, the correlations are readily seen to exhibit crossover from weak coupling behavior, $\langle \sigma_z \sigma_z \rangle \sim \text{const}$, to the non-Fermi liquid behavior of Eq(42) for $\tau \gtrsim \Delta_K^{-2} \tau_c$.

The 2CK fixed point, formally, corresponds to $\Delta_K \rightarrow \infty$. It is more convenient, however, to allow instead the normalization of γ to change, corresponding to letting the coefficient of the $\int \gamma \partial_\tau \gamma d\tau$ in the Lagrangian vary. At the fixed point, this coefficient, say g_γ , will be zero, while Δ_K becomes a constant; the correlations of γ are then simply the inverse (in frequency space) of those of β_I , i.e. a pure power law Eq(B10). To connect the two regimes together, one could choose to renormalize so that, for example, $\frac{\Delta_K^2}{4\pi} + g_\gamma = 1$ [a particularly convenient choice; (see Eq(B12))], by rescaling γ under renormalization by a Δ_K -dependent amount. Details about this procedure are given in Appendix B. [Note, however, that the resulting fixed point Hamiltonian (with $\Delta_K = \sqrt{4\pi}$) will *not* have the pseudo-spin $SU(2)$ symmetry of the pure 2CK problem. This is because the RG approach we have implemented here, including the unitary transformation of Eq(28), is inherently anisotropic in channels (equivalent to spin of conventional 2CK problem). In Appendix C we will show that this anisotropy will not affect the results.]

At the 2CK fixed point, the above scaling implies $\gamma \sim \tau^{-\frac{1}{2}}$, while δ is still dimensionless so that the RG eigenvalues of the other operators can be read off immediately: $+1/2$ for Δ_+ , the unique relevant operator consistent with the symmetries of the problem; $-1/2$ for $\bar{q}$, $\bar{y}$ and w , which are the three leading irrelevant operators discussed so far (the fourth will appear in Appendix C); and 0 for u , which is marginal but redundant, in that it does not affect the impurity dynamics.

The operator corresponding to $\bar{q}$ does not give rise to any terms which couple δ linearly to the α 's and β 's and hence will not generate Δ_+ . It is the leading irrelevant operator for the 2CK problem identified by Sengupta and Georges¹⁰. An artificial symmetry is responsible for its special role, indeed just the one discussed earlier: Eq(35), $\Phi_c \rightarrow \pi - \Phi_c$ corresponds to the discrete symmetry $\alpha_c \rightarrow -\alpha_c$ and $\beta_X \rightarrow -\beta_X$ which does not have a natural representation even in terms of Φ_c and Φ_s . But if present, this artificial symmetry excludes the generation of terms like Δ_+ , $\bar{y}$ and w , even in the presence of $\bar{q}$. In fact, without these terms, the extra artificial symmetry is really an $O(2)$ symmetry in the $\{\alpha_c, \beta_X\}$ pair, consisting of a $U(1)$ of rotations in the (α_c, β_X) plane, combined (in a non-commutative way) with Z_2 , the usual site interchange symmetry, which takes $\alpha_c \rightarrow -\alpha_c$ and $\beta_X \rightarrow \beta_X$. This is exactly analogous to the $O(2)$ symmetry present in the model treated by Emery and Kivelson¹⁸ and Sengupta and Georges¹⁰.

In contrast, the irrelevant operators w and $\bar{y}$, break the artificial symmetry and yield, at lowest order, the generation of the relevant operator Δ_+ (see Eq(B14))

$$\frac{d\Delta_+}{dl} = \frac{1}{2}\Delta_+ + 2\bar{y}\Delta_K + \frac{wu}{\sqrt{2}(1+u^2)}, \quad (43)$$

consistent with expectations from weak coupling. This implies as stated earlier, that the critical point will not be exactly at $\Delta_+ = 0$.

Of the three leading irrelevant operators, it should be noted that, although $\bar{y}$ has the same scaling dimension as $\bar{q}$ and w , it has a different role close to the 2CK fixed point. The reason is that it couples to the term $i\gamma\beta_I$, already present at the fixed point. But the Δ_K term at the fixed point suppresses fluctuations of β_I , causing the leading term in the correlations of β_I to *vanish* at the fixed point (with sub-dominant terms caused by $g_\gamma \neq 0$). As a result, unlike $\bar{q}$ and w which each yield $O(T \ln T)$ contributions to the impurity specific heat¹⁰ — a key feature of a 2CK non-Fermi liquid — the singular part arising from $\bar{y}$ is only of $O(T^3 \ln T)$, for temperatures $T \ll T_K$. Thus only *two* of the above independent leading irrelevant operators give leading singular specific heat corrections. In fact, there is a third one, involving only spin degrees of freedom, which is discussed in Appendix C.

In Section VI, the conditions for accessibility of the 2CK fixed point are analyzed. We turn here to further analysis of the behavior near the 2CK fixed point.

D. Symmetry Breaking Operators

Up to this point we have only dealt with an electron-impurity system which is invariant under Z_2 ($1 \leftrightarrow 2$ interchange) and spin $SU(2)$ symmetry. However, in a realistic situation of an impurity in a metal, there will generally be a non-zero, although possibly small asymmetry between the two sites which will break the Z_2 symmetry. Furthermore, in the presence of a magnetic field, the equivalence between the two spin channels will be lost, leading to a situation similar to the anisotropic Kondo problem^{1,22,23}. As might be expected, in both cases, the symmetry breaking terms are relevant and in their presence the system flows away from the 2CK fixed point. In this section, we will briefly comment on the effects of symmetry breaking terms close to the 2CK fixed point.

It is clear from the discussion in the previous section that for an operator to be relevant close to the 2CK fixed point, it has to be of the form $i\delta\chi$ where χ is a Majorana fermion of scaling dimension $1/2$. From the ten Majorana fermions (four pairs of α_μ, β_μ and γ, δ , all listed in Table I) we can make nine such operators. Excluding $i\delta\alpha_{ec}$ and $i\delta\beta_{ec}$ due to total electron number conservation (which only allows α_{ec} and β_{ec} to appear together as $\alpha_{ec}\beta_{ec}$) we are left with seven possible terms.

From the transformation properties under the discrete symmetries of the system, listed in Table I, it can be seen that β_c and β_s have the same symmetries; indeed this is why the Δ_1 and Δ_- terms in Eq(38) could be combined

into the Δ_K term of Eq(40). In the presence of small $i\delta\beta_c$ and $i\delta\beta_s$ terms, a small rotation of the (δ, γ) pair as well as a small additional rotation of the (β_c, β_s) pair can be performed to yield just a slightly modified $i\Delta_K\gamma\beta_I$ term, and a single remaining relevant perturbation, the $i\delta\beta_X$ term. The extra operators ($i\gamma\beta_X$ and $i\delta\beta_I$) are thus “redundant”.²⁴ Therefore at the 2CK fixed point, there are exactly *six relevant* operators, all with RG eigenvalue of 1/2, like Δ_+ . These, along with their symmetry properties, are listed in Table II.

The first three operators in Table II do not break the spin $SU(2)$ symmetry. Among these, the first, our familiar $i\Delta_+\delta\alpha_c$, is interchange and time-reversal symmetric, corresponding to the relevant part of the channel pseudo-spin operator S_x . Correspondingly, the second, $i\delta\gamma$, breaks interchange symmetry but is time reversal invariant, corresponding to a S_z operator. This will result from simple asymmetry between the impurity energies of the two sites, i.e. a σ_z term in the original Hamiltonian. Finally, $i\delta\beta_c$ (or, equivalently, $i\delta\beta_s$) breaks interchange and time reversal which makes it an imaginary operator, i.e. it is generated by a S_y operator in the channel sector, corresponding to complex hopping matrix elements.

The relevant spin $SU(2)$ breaking operators are $i\delta\alpha_s$, $i\delta\alpha_{es}$ and $i\delta\beta_{es}$; these correspond to joint electron-impurity hops accompanied by a spin flip or carrying electronic spin. They correspond to combinations of $\sigma_+ (c_{1\uparrow}^\dagger c_{2\uparrow} - c_{1\downarrow}^\dagger c_{2\downarrow})$, $\sigma_+ c_{1\downarrow}^\dagger c_{2\uparrow}$, $\sigma_+ c_{1\uparrow}^\dagger c_{2\downarrow}$ and their hermitian conjugates and are discussed in Appendix B (see Eq(B16) and Eq(B17)). The first corresponds to the “flavor” anisotropy term in the conventional two channel Kondo model^{1,22,23}, and, being interchange and time reversal symmetric but odd under spin flip, is induced by a magnetic field in the z -direction. Interestingly, the remaining two relevant spin $SU(2)$ breaking operators are *odd* under the Z_2 interchange transformation. This means that in order to adjust them to zero in a non-zero external magnetic field, one would have to tune also terms that break the interchange symmetry of the problem, making any additional novel, finite magnetic field, non-Fermi liquid fixed points (analogous to that found in zero field for $\Delta_+ \gg \Delta_1, \Delta_-$), extremely hard to observe in a system that does not have interchange symmetry.

V. ADDITIONAL INTERMEDIATE COUPLING FIXED POINT

In the previous section we analyzed the behavior of the system close to the 2CK fixed point, that corresponds to the limit $\Delta_K \gg \Delta_+$. There, we showed that γ , “half” of the impurity, acquired non-trivial dynamics, which essentially gave rise to the non-Fermi liquid behavior of the system. However, as is evident by examining Eq(40), it should be, in principle, possible to get the same type of non-Fermi behavior if the inequality above were reversed ($\Delta_K \ll \Delta_+$).

Indeed, following the same arguments analyzed above, we see that when $\Delta_K = \bar{q} = \bar{y} = w = u = 0$ and $\Delta_+ \neq 0$ we have a critical point, which has an extra artificial $U(1)$ symmetry, namely rotations in the (β_I, β_X) plane. At this secondary fixed point, δ , the other “half” of the impurity, acquires dynamics, rather than γ . The operators $\bar{q}$, $\bar{y}$, $i\gamma\beta_I\Psi_{ec}^\dagger\Psi_{ec}$ and $i\gamma\beta_X\Psi_{ec}^\dagger\Psi_{ec}$ have scaling dimension 3/2 and thus are irrelevant with RG eigenvalue -1/2. But of these, only the last two will give singular specific heat corrections ($O(T \ln T)$).

However, there is an important difference from the primary 2CK fixed point discussed earlier. About this fixed point, there are *two* relevant operators, consistent with the symmetries of the model, namely $i\gamma\beta_I$ and $i\gamma\beta_X$ (or, equivalently, $i\gamma\beta_c$ and $i\gamma\beta_s$), both with dimension 1/2. These correspond simply to Δ_1 and Δ_- before the change of variables (Eq(B6)) leading to Eq(40); they can be generated from nonzero q and y . The existence of two relevant operators is due to the fact that β_I and β_X transform the *same* way under the discrete symmetries; thus the artificial $U(1)$ symmetry *cannot* be extended into an $O(2)$ group (as was the case for the 2CK fixed point). As a result, this new fixed point is harder to find in the interchange symmetric case than the primary 2CK fixed point, as it requires the impurity-single-electron hopping term to be small and the two-electron-plus-impurity and the simple impurity hopping terms to be almost exactly equal at the Kondo scale. It should be noted, however, that the *total* number of relevant, dimension 1/2, operators around this fixed point is again six, including the above mentioned ones, together with $i\gamma\delta$ and the three spin $SU(2)$ symmetry breaking operators, $i\gamma\alpha_s$, $i\gamma\alpha_{es}$ and $i\gamma\beta_{es}$, as discussed in Section IV.E and listed in Table II, indicating that the fixed point symmetry is again that of the conventional 2CK model. This is supported by the fact that, just like in the case of the 2CK fixed point, there are four operators with scaling dimension 3/2, albeit with completely different symmetries.

Finally, some comments are needed on the the nature of this novel fixed point. We should first stress the *absence* of Δ_1 , the impurity-single-electron hopping term, which together with the Q_0 term (see Eq(8)) would form the conventional Kondo-like interaction term. Hence, the appearance of non-Fermi liquid behavior does *not* originate from the competition between the spin up and spin down electrons to form a channel-pseudo-spin singlet ground state with the impurity, but, rather, in the presence of strong impurity-electron repulsion ($Q_0 \approx 1/2$), from the competition between bare impurity tunnelling and two-electron-plus-impurity tunnelling. Formally, there is an analogy with the conventional two channel flavor-anisotropic Kondo model,²² with $\Delta_\pm$ playing the role of the Kondo couplings of the two “flavor” channels, which can be seen in the left hand side of Eq(30) if Φ_c is substituted by Φ_s . When $\Delta_0 \neq \Delta_2$ one flavor channel couples more strongly to the impurity therefore screening it alone at low energy, which results in usual Fermi liquid

behavior. However, if $\Delta_0 = \Delta_2$ the flavor anisotropy disappears and the system flows to a non-Fermi liquid fixed point (provided Δ_1 is zero). As a result, although this fixed point may be in the same universality class as the conventional 2CK model, the mechanism that brings it about is completely different physically.

VI. ACCESSIBILITY OF THE TWO CHANNEL KONDO FIXED POINT AND CONCLUSIONS

In the previous sections we have shown how the physical operators in the tunnelling impurity problem behave near the two channel Kondo fixed point. In particular, we observed that a linear combination, $\Delta_+ = \Delta_0 + \Delta_2$, of the bare impurity hopping, Δ_0 , and the impurity-plus-two-electron hopping, Δ_2 , is relevant and drives the system away from this special critical point resulting in conventional Fermi liquid behavior at low temperatures, as for the usual one-channel Kondo system. If one could somehow tune Δ_+ , (or one other coupling such as the electronic hopping y) then one might be able to tune through the critical point and find the 2CK non-Fermi liquid behavior at low temperatures in the vicinity of a critical coupling. Unfortunately, such tuning over an adequate range is probably difficult to achieve. Thus one probably has to rely on the hope that a natural regime of couplings will lead to flow under renormalization close to the 2CK fixed point. Vladár and Zawadowski³ appear to suggest that this should be the case. Unfortunately, more complete analysis implies the converse, that only for fortuitous reasons would the impurity system — even without asymmetry between the sites — exhibit 2CK behavior at low T .

In this last section, we use the weak coupling analysis of Section III and Appendix A combined with the intermediate coupling analysis of Section IV and Appendix B, to find criteria for approaching close to the 2CK fixed point.

Since we are interested in systems in which the Kondo temperature is much less than the bandwidth, the weak hopping behavior will control the relative strengths of couplings at the Kondo scale, at which the first of the hopping terms becomes of order the renormalized bandwidth. Operators which are irrelevant for weak hopping will flow away rapidly under renormalization, changing by finite amounts the remaining parameters, q , y and the $\{\Delta_i\}$, $i = 0, 1, 2$. For example, the complicated hopping-scattering term, w , which was discussed in Section IV and plays a role near the intermediate coupling fixed point, will be of order the hopping terms $\{\Delta_i\}$ or smaller initially and flow away rapidly, modifying, among other terms, Δ_+ as in Eq(43), in the process. This will be the main role of such a term and we can incorporate its effects into a modified “bare” Δ_0 and Δ_2 . We thus start at an energy scale substantially below the bandwidth at which the important parameters for $Q \in [0, 1]$ are just

q , y and the $\{\Delta_i\}$ at this scale, the irrelevant operators having become small. The relevant eigenvalues for the hopping about the zero hopping fixed manifold will be universal. For small y they are given by Eq(24).

If Q is initially small, i.e. $q \leq 1/2$, λ_+ will be substantially larger than the other eigenvalues and thus a particular linear combination of the $\{\Delta_i\}$ will grow fast. Unfortunately, as can be seen from Appendix A, this combination ($\tilde{\Delta}_0$) is the wrong one to yield flow near the 2CK fixed point as it *includes* Δ_+ . Only if this combination, $\tilde{\Delta}_0$, is initially very small relative to a power of another linear combination of the same $\{\Delta_i\}$, can behavior near the 2CK critical point be obtained; furthermore the criteria become more stringent the lower the Kondo temperature, as shown in Appendix A.

Better prospects occur when $Q \approx 1/2$ (i.e. q small). Unfortunately, even if the one-electron plus impurity hopping term, Δ_1 , were initially much bigger than Δ_0 and Δ_2 , the purely electronic hopping term y — determined basically by the spatial separation of the two impurity sites¹⁴ — would combine with Δ_1 to generate the *wrong* combination, Δ_+ , of Δ_0 and Δ_2 , as in Eq(43). Thus, again, unless y is small the criteria from Appendix A are very strict, as could be anticipated from the y dependence of $\lambda_{\pm}$.

The best prospects are thus for q and y both small so that the eigenvalues are all comparable. But this is just the condition for the analysis of Section IV and Appendix B via refermionization to be valid. To get near the 2CK fixed point, Δ_+ must remain small, thus we can study the RG equations Eq(B14) to leading order in Δ_+ and the linear combinations of y and q , $\bar{y}$ and $\bar{q}$;

$$\begin{aligned} \frac{d\Delta_K}{dl} &= \frac{\Delta_K}{2} \left(1 - \frac{\Delta_K^2}{4\pi} \right) \\ \frac{d\bar{q}}{dl} &= -\frac{\Delta_K^2}{8\pi} \bar{q} \\ \frac{d\bar{y}}{dl} &= -\frac{\Delta_K^2}{8\pi} \bar{y} \end{aligned} \quad (44)$$

and, ignoring w from Eq(43)

$$\frac{d\Delta_+}{dl} = \frac{1}{2}\Delta_+ + 2\bar{y}\Delta_K. \quad (45)$$

At the intermediate coupling 2CK fixed point, the convention we have chosen yields $\Delta_K^* = \sqrt{4\pi}$, so that $\bar{y}$ and $\bar{q}$ have the correct eigenvalues there as well as for weak coupling ($\Delta_K \approx 0$).

Integrating the above equations we find that the criterion to flow to the 2CK fixed point is, to leading order in Δ_+ and $\bar{y}$

$$\Delta_+ - \frac{2\bar{y}\Delta_K}{1 - \frac{\Delta_K^2}{4\pi}} \ln \left| \frac{\Delta_K^2}{4\pi} \right| = 0 \quad (46)$$

The parameters can be evaluated at any scale where the intermediate coupling RG equations are valid, i.e. for small Δ_+ , $\bar{y}$ and $\bar{q}$. If these are small at the starting

scale, then their starting values can be used. Note that since $\Delta_K = \sqrt{\Delta_1^2 + (\Delta_0 - \Delta_2)^2}$, unless Δ_0 and Δ_2 are almost exactly equal and opposite, Δ_K will be at least as big as Δ_+ initially so that $|\bar{y} \ln \Delta_K|$ needs to be small rather than just y , even in the best case of $q = 1/2 - Q$ small (recall that $\bar{y}$ is a linear combination of q and y given by Eq(B7)).

We thus see that the flows near the 2CK fixed point, in particular the generation via Eq(43) of the unique relevant operator, Δ_+ from other operators, yield stringent conditions for the accessibility of the non-Fermi liquid 2CK behavior. If there is no symmetry between the two sites, then the presence of a second relevant operator (see Section IV) makes prospects even worse.

This work strongly suggests that to observe two-channel-Kondo-like non-Fermi liquid behavior of an impurity hopping between two sites in a metal one must either be able to tune some parameters over a substan-

tial range, or be extremely lucky. This casts doubt on the interpretation of Ralph *et al*^{4,5} of their narrow constriction tunnelling data. One possibility, although perhaps farfetched, is that these might be some kind of defects tunnelling in environments with higher symmetry, or at least approximate symmetry. In another paper, we will show how an impurity hopping among *three* sites with triangular symmetry can, without fine tuning, lead to a two channel Kondo behavior at low temperatures.

ACKNOWLEDGMENTS

We would like to thank Andreas Ludwig, Jinwu Ye, Dan Ralph, Jan von Delft, Igor Smolyarenko and especially Anirvan Sengupta for useful discussions. This work was supported by the National Science Foundation via grant DMR 91-06237.

APPENDIX A: DETAILS OF WEAK-INTERMEDIATE COUPLING CROSSOVER

In this Appendix we analyze the dependence of the crossover energy scale, T_K , and other quantities as a function of initial values of various coupling constants. For simplicity, we ignore the coupling to the even parts of the electron fields (and hence terms like u and w in Eq(29)). Since in the weak coupling RG flow equations (Eq(23)) of Δ_0 , Δ_1 and Δ_2 , q and y are essentially constant, these equations can be trivially integrated. The solutions for the two quantities of interest at the intermediate coupling region, i.e. $\Delta_+(l)$ and $\Delta_K(l)$ are

$$\Delta_+(l) = \Delta_0(l) + \Delta_2(l) = \tilde{\Delta}_0 e^{\lambda_+ l} + \tilde{\Delta}_2 e^{\lambda_- l} \quad (A1)$$

and

$$\Delta_K(l) = \sqrt{\Delta_1^2(l) + \Delta_-^2(l)} = \sqrt{(\tilde{\Delta}_1 e^{\lambda_0 l})^2 + (\tilde{\Delta}_0 e^{\lambda_+ l} - \tilde{\Delta}_2 e^{\lambda_- l})^2}, \quad (A2)$$

with the weak coupling eigenvalues given by Eq(24)

$$\begin{aligned} \lambda_{\pm} &= \frac{1}{2} - 2q_0^2 \pm 2\sqrt{q_0^2 + y_0^2} \\ \lambda_0 &= \frac{1}{2} - 2q_0^2 \end{aligned}$$

Note that the weak coupling eigenvalues for the relevant hopping terms should be universal.¹⁷ But since we have ignored higher order terms in y , (as well as renormalization of q from high energy scales which make it related to a phase shift rather than a coupling constant) the expressions for the eigenvalues in terms of the original y_0 and q_0 will not be exact.

The couplings that appear in Eqs(A1) and (A2), $\tilde{\Delta}_0$, $\tilde{\Delta}_1$ and $\tilde{\Delta}_2$, are linear combinations of the original Δ_0 , Δ_1 and Δ_2 :

$$\begin{aligned} \tilde{\Delta}_0 &= \frac{1}{2\sqrt{q^2 + y^2}} \left\{ \left(\sqrt{q^2 + y^2} + q \right) \Delta_0 + \left(\sqrt{q^2 + y^2} - q \right) \Delta_2 + y \Delta_1 \right\} \\ \tilde{\Delta}_1 &= \frac{1}{\sqrt{q^2 + y^2}} \{ y (\Delta_2 - \Delta_0) + q \Delta_1 \} \\ \tilde{\Delta}_2 &= \frac{1}{2\sqrt{q^2 + y^2}} \left\{ \left(\sqrt{q^2 + y^2} - q \right) \Delta_0 + \left(\sqrt{q^2 + y^2} + q \right) \Delta_2 - y \Delta_1 \right\}. \end{aligned} \quad (A3)$$

The weak coupling flow equations in Eq(23) break down at the crossover scale $T_K = \tau_c^{-1} e^{-l^*}$, which is reached when the first of $\Delta_+(l^*)$ and $\Delta_K(l^*)$ becomes of order unity.

At this scale, $\bar{q}$ and $\bar{y}$, defined in Eq(B7) and appearing in Eq(40) are

$$\begin{aligned}\bar{q} &= \sqrt{q^2 + y^2} \frac{\tilde{\Delta}_1}{(T_K \tau_c)^{\lambda_0}} \\ \bar{y} &= \sqrt{q^2 + y^2} \left(\frac{\tilde{\Delta}_0}{(T_K \tau_c)^{\lambda_+}} - \frac{\tilde{\Delta}_2}{(T_K \tau_c)^{\lambda_-}} \right).\end{aligned}\tag{A4}$$

In order to flow close to the 2CK fixed point, it is necessary that $\Delta_K(l^*) = O(1)$ first, with $\Delta_+(l^*) \ll 1$. There are two regions of parameter space, in terms of $\tilde{\Delta}_0$, $\tilde{\Delta}_1$ and $\tilde{\Delta}_2$, where this is realizable. The first is

$$\begin{aligned}\frac{\left| \tilde{\Delta}_0 \right|^{\lambda_0}}{\left| \tilde{\Delta}_1 \right|^{\lambda_+}} &\ll 1 \\ \frac{\left| \tilde{\Delta}_2 \right|^{\lambda_0}}{\left| \tilde{\Delta}_1 \right|^{\lambda_-}} &\ll 1,\end{aligned}\tag{A5}$$

in which case the crossover scale is

$$T_K \approx \left| \tilde{\Delta}_1 \right|^{\frac{1}{\lambda_0}} \tau_c^{-1}.\tag{A6}$$

The other region corresponds to the second square under the square-root sign in Eq(A2) being of order unity. In this case the inequality of Eq(A5) is reversed, $\tilde{\Delta}_0$ and $\tilde{\Delta}_2$ have to be opposite in sign, ($\tilde{\Delta}_0 \tilde{\Delta}_2 < 0$) and

$$\left| \frac{\left| \tilde{\Delta}_2 \right|^{\lambda_+}}{\left| \tilde{\Delta}_0 \right|^{\lambda_-}} - 1 \right| \ll 1.\tag{A7}$$

Then T_K is

$$T_K \approx \left| \tilde{\Delta}_0 \right|^{\frac{1}{\lambda_+}} \tau_c^{-1} \approx \left| \tilde{\Delta}_2 \right|^{\frac{1}{\lambda_-}} \tau_c^{-1}.\tag{A8}$$

The opposite limit corresponds to $\tilde{\Delta}_+(l^*)$ becoming of order unity first. This will happen when Eqs(A7)-(A8) hold, but with $\tilde{\Delta}_0$ and $\tilde{\Delta}_2$ having the same sign ($\tilde{\Delta}_0 \tilde{\Delta}_2 > 0$).

It can be seen from Eq(A3) that the above two conditions Eq(A5) and Eq(A7) are complicated functions of the initial parameters q , y and the Δ 's. In the remainder of this Appendix we will analyze two limiting cases in terms q and y .

The first limit is $|q| \gg |y|$. From Eq(A3) we see that for $q > 0$ this corresponds to $\tilde{\Delta}_i \approx \Delta_i$ for $i = 0, 1$ and 2 , in which case the conditions in terms of Δ_i are read directly from Eqs(A5)-(A7). It should be noted that small initial $Q_0 = \frac{1}{2} - q_0$, i.e. weak coupling of the electrons to the impurity position, (Eq(8)), lies in this regime.

In the second limiting case, whence $|q| \ll |y|$,

$$\begin{aligned}\tilde{\Delta}_0 &\approx \frac{1}{2} (\Delta_0 + \Delta_2 + \text{sgn}(y) \Delta_1) \\ \tilde{\Delta}_2 &\approx \frac{1}{2} (\Delta_0 + \Delta_2 - \text{sgn}(y) \Delta_1) \\ \left| \tilde{\Delta}_1 \right| &\approx \left| \Delta_2 - \Delta_0 \right|.\end{aligned}\tag{A9}$$

In this region, Eq(A5) becomes, for $y > 0$, approximately,

$$\begin{aligned}(\Delta_0 + \Delta_2) + \Delta_1 &\ll |\Delta_0 - \Delta_2|^{1+4y} \\ (\Delta_0 + \Delta_2) - \Delta_1 &\ll |\Delta_0 - \Delta_2|^{1-4y},\end{aligned}\tag{A10}$$

while Eq(A7), for $y > 0$, corresponds to

$$\begin{aligned} |\Delta_1| &> |\Delta_0 + \Delta_2| \\ |(\Delta_0 + \Delta_2) - \Delta_1|^{1-4y} &\approx |(\Delta_0 + \Delta_2) + \Delta_1|^{1+4y}. \end{aligned} \quad (\text{A11})$$

In conclusion, for both limiting situations, as well as for the intermediate region, where $|q| \approx |y|$, there is a finite, albeit small, region of initial parameters where the approach to the 2CK fixed point at low energies is indeed possible.

Finally, we demonstrate that the weak coupling condition for flowing to the 2CK fixed point, Δ_+ (l^*) small, evaluated at the Kondo temperature, $T_K = \tau_c^{-1} e^{-l^*}$, matches with the one derived from the intermediate coupling side (Eq(46)) in the parameter region where both are valid, i.e. small initial $\bar{q}$, $\bar{y}$, Δ_+ and Δ_K . To linear order in q , y and Δ_+ we have from Eq(A2), $\Delta_K(l) \approx e^{l/2} \Delta_K$ and hence

$$T_K \approx \Delta_K^2 \tau_c^{-1}. \quad (\text{A12})$$

The condition for criticality becomes from Eq(A1)

$$\frac{\tilde{\Delta}_0}{(T_K \tau_c)^{\lambda_+}} + \frac{\tilde{\Delta}_2}{(T_K \tau_c)^{\lambda_-}} = 0. \quad (\text{A13})$$

After expanding the factors $T_K^{-\lambda_{\pm}}$ to leading order in q and y and writing $l^* \approx -\ln(\Delta_K^2)$ this yields

$$\left(\frac{\tilde{\Delta}_0}{(T_K \tau_c)^{\lambda_0}} + \frac{\tilde{\Delta}_2}{(T_K \tau_c)^{\lambda_0}} \right) - 2\sqrt{q^2 + y^2} \left(\frac{\tilde{\Delta}_0}{(T_K \tau_c)^{\lambda_0}} - \frac{\tilde{\Delta}_2}{(T_K \tau_c)^{\lambda_0}} \right) \ln \Delta_K^2 = 0. \quad (\text{A14})$$

Noting that the terms in the first parentheses is just Δ_+ evaluated at T_K , to lowest order in q and y , while the term multiplying the logarithm is just $2\bar{y}$ at T_K , to lowest order in q and y (see Eq(A4)), we recover the condition Eq(46) derived from the refermionization calculation, for the initial Δ_K small.

APPENDIX B: DETAILED ANALYSIS OF INTERMEDIATE COUPLING POINT

In this Appendix the details of the intermediate coupling point analysis are discussed. The starting point is the bosonization of the Hamiltonian in the Eqs(7-9), using Eq(10) and Eq(11), but inserting and keeping track of ordering factors which ensure proper anticommutation relations between different fields. The original fermions $c_{1\uparrow}$, $c_{2\uparrow}$, $c_{1\downarrow}$, $c_{2\downarrow}$ are bosonized in the following way (rather than just the simple form of Eq(10)):

$$\begin{aligned} c_{1\uparrow} &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_{1\uparrow}} \\ c_{2\uparrow} &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_{2\uparrow}} e^{i\pi(N_{1\uparrow} + N_{2\uparrow})} \\ c_{1\downarrow} &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_{1\downarrow}} e^{i\pi(N_{1\uparrow} + N_{2\uparrow})} \\ c_{2\downarrow} &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_{2\downarrow}} e^{i\pi(N_{1\uparrow} + N_{2\uparrow} + N_{1\downarrow} + N_{2\downarrow})} \end{aligned} \quad (\text{B1})$$

where $N_{1\uparrow}$, $N_{2\uparrow}$ etc. are the number operators for the associated fields:

$$N_{1\uparrow} = \int dx c_{1\uparrow}^\dagger(x) c_{1\uparrow}(x) = \frac{1}{2\pi} \int d\epsilon c_{1\uparrow\epsilon}^\dagger c_{1\uparrow\epsilon} \quad (\text{B2})$$

etc. with the total number of electron fields $N_{tot} = N_{1\uparrow} + N_{2\uparrow} + N_{1\downarrow} + N_{2\downarrow}$ being always conserved. Note that since the number operators have integer eigenvalues the exponential factors in Eq(B1) can only take the values plus and minus one. With these extra ordering factors the only change in the effective Hamiltonian of Eq(13) is the appearance of

$$\exp[i\pi(N_{1\uparrow} + N_{2\uparrow})] \quad (\text{B3})$$

multiplying y and Δ_1 . Since the combination $N_{1\uparrow} + N_{2\uparrow}$ is conserved in the absence of spin-flip processes, the exponential factor can be set to plus or minus unity and be disregarded. This can be seen more clearly if, using the number operators N_c , N_s , N_{ec} and N_{es} defined in an analogous way as the corresponding Φ -fields in Eq(19) and Eq(20), we realize that $N_{1\uparrow} + N_{2\uparrow} = N_{ec} + N_{es}$. Since $N_{tot} = 2N_{ec}$ which is always conserved, the exponential factor of Eq(B3) will be conserved if N_{es} is constant, i.e. if no spin-flip processes can appear. [Note that in the absence of spin flip processes, one can always choose a convention that different spin species *commute* rather than anti-commute.]

After performing the unitary transformation using $U = \exp[-\frac{i}{2}\sigma_z\Phi_c]$ we get the transformed Hamiltonian $\tilde{\mathcal{H}}$ of Eq(29). Next, following Emery and Kivelson¹⁸, we refermionize the exponentials of the Bose fields Φ_μ with $\mu = c, s, ec, es$ and also fermionizing σ_- keeping track of ordering operators, as in Eq(B1), to ensure proper anticommutation relations between different fermion fields $\{\Psi_\mu\}$ and d :

$$\begin{aligned}\Psi_s &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_s} e^{i\pi N_s} \\ \Psi_c &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_c} e^{i\pi(N_{ec}+N_{es}+N_s+N_d)} \\ d &= \sigma_- e^{i\pi(N_s+N_{es}+N_{ec})} \\ \Psi_{es} &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_{es}} e^{i\pi(N_{es}+N_{ec}+N_s)} \\ \Psi_{ec} &= \frac{1}{\sqrt{2\pi\tau_c}} e^{i\Phi_{ec}} e^{i\pi N_s}\end{aligned}\tag{B4}$$

where N_μ is the number operator for the μ -fermions and $N_d = d^\dagger d$. The Hamiltonian now becomes

$$\begin{aligned}\hat{\mathcal{H}} &= \mathcal{H}_0 + \frac{1}{2\sqrt{2\pi\tau_c}} \{ -\Delta_1 (d + d^\dagger) (\Psi_s - \Psi_s^\dagger) + \Delta_- (d + d^\dagger) (\Psi_c - \Psi_c^\dagger) \} \\ &\quad + 2u\Psi_{ec}^\dagger \Psi_{ec} + \frac{\Delta_+}{2\sqrt{2\pi\tau_c}} (d - d^\dagger) (\Psi_c + \Psi_c^\dagger) \\ &\quad + \pi y (2d^\dagger d - 1) (\Psi_c + \Psi_c^\dagger) (\Psi_s - \Psi_s^\dagger) \\ &\quad - 2\pi q \Psi_c^\dagger \Psi_c (2d^\dagger d - 1) - \frac{\sqrt{2\pi\tau_c}w}{2} (d - d^\dagger) (\Psi_c + \Psi_c^\dagger) \Psi_{ec}^\dagger \Psi_{ec}.\end{aligned}\tag{B5}$$

Note that we have chosen the ordering operators in the refermionization, so that the ordering operators appearing in the bosonized Hamiltonian cancel.

To simplify the Hamiltonian we introduce a set of eight Majorana fermions of Eq(36), so that the Hamiltonian takes the form of Eq(38). Their transformation properties under the various symmetries are listed in Table I. Now, as it is clear from this Table, β_s and β_c have the same transformation properties under the discrete symmetries of the system. Therefore, we can make linear combinations of the two so that only one will couple to the γ Majorana fermion. Defining

$$\begin{aligned}\beta_I &= \frac{\Delta_- \beta_c - \Delta_1 \beta_s}{\Delta_K} \\ \beta_X &= \frac{\Delta_1 \beta_c + \Delta_- \beta_s}{\Delta_K}\end{aligned}\tag{B6}$$

and

$$\begin{aligned}\bar{q} &= \frac{q\Delta_1 - y\Delta_-}{\Delta_K} \\ \bar{y} &= \frac{q\Delta_- + y\Delta_1}{\Delta_K}\end{aligned}\tag{B7}$$

where $\Delta_K = \sqrt{\Delta_1^2 + \Delta_-^2}$ as in Eq(39), we arrive to the Hamiltonian in the form of Eq(40)

$$\begin{aligned}\hat{\mathcal{H}} &= \mathcal{H}_0 + 2iu\alpha_{ec}\beta_{ec} + \frac{i}{\sqrt{2\pi\tau_c}} (\Delta_K \gamma \beta_I + \Delta_+ \delta \alpha_c) \\ &\quad + 4\pi \bar{q} \gamma \delta \alpha_c \beta_X + 4\pi \bar{y} \gamma \delta \alpha_c \beta_I + \sqrt{2\pi\tau_c} w \delta \alpha_c \alpha_{ec} \beta_{ec},\end{aligned}$$

In the RG the operator $i\gamma\beta_X$ is *redundant*; under renormalization it can be generated but can always be rotated away.²⁴

Before deriving the RG flow equations for the various operator amplitudes in Eq(40) close to the intermediate coupling fixed point, we should first locate the fixed point. As discussed in Section IV, formally it should lie at $\Delta_K = \infty$. This can be seen in the following way. At the fixed point, with $\Delta_+ = \bar{q} = \bar{y} = w = 0$ for convenience, the Δ_K term in Eq(40) should have the same scaling dimension as $\mathcal{H}_0$, i.e. unity. Defining the Fourier transform of the non-interacting ($\Delta_K = 0$), imaginary time Green's functions for γ and β_I as

$$G_{0\gamma}(i\omega) = \frac{1}{i\omega g_\gamma} = -F.T. \langle T_\tau [\gamma(\tau) \gamma(0)] \rangle_0 \quad (B8)$$

and

$$G_{0\beta_I}(i\omega) = -i \frac{\text{sgn}(\omega)}{2} = -F.T. \langle T_\tau [\beta_I(\tau) \beta_I(0)] \rangle_0, \quad (B9)$$

respectively, where g_γ is related to the normalization of γ , as discussed in Section IV, and $F.T.$ stands for Fourier transform, we see that the corresponding interacting ($\Delta_K \neq 0$) Green's functions are

$$G_\gamma(i\omega) = \frac{1}{G_{0\gamma}^{-1}(i\omega) - \frac{\Delta_K^2}{2\pi\tau_c} G_{0\beta_I}(i\omega)} = \frac{1}{ig_\gamma\omega + \frac{i\Delta_K^2 \text{sgn}(\omega)}{4\pi\tau_c}} \quad (B10)$$

and

$$G_{\beta_I}(i\omega) = \frac{1}{G_{0\beta_I}^{-1}(i\omega) - \frac{\Delta_K^2}{2\pi\tau_c} G_{0\gamma}(i\omega)} = \frac{1}{2i\text{sgn}(\omega) + \frac{i\Delta_K^2}{2\pi\tau_c g_\gamma\omega}} \quad (B11)$$

This means that the naive scaling dimensions of γ and β_I ($\omega \rightarrow 0$) with Δ_K and g_γ *finite* are 1/2 and 3/2, respectively, adding to 2. Thus some combination of Δ_K and g_γ must be infinite at the fixed point. But more useful than letting $\Delta_K \rightarrow \infty$ is to keep Δ_K finite and let $g_\gamma \rightarrow 0$. A convenient form, which we will adopt, is to choose the normalization of γ so that

$$g_\gamma + \frac{\Delta_K^2}{4\pi} = 1 \quad (B12)$$

This fixes the scaling dimension of γ to be 1/2 at the fixed point, but also correctly gives the conventional scaling as $\Delta_K \rightarrow 0$. This choice determines the part of RG flow equations for Δ_K and g_γ that does not involve any other parameters:

$$\begin{aligned} \left. \frac{d\Delta_K}{dl} \right|_{\text{rescaling}} &= \frac{g_\gamma}{2} \Delta_K = \frac{\Delta_K}{2} - \frac{\Delta_K^3}{8\pi} \\ \left. \frac{dg_\gamma}{dl} \right|_{\text{rescaling}} &= -\frac{\Delta_K^2}{4\pi} g_\gamma \end{aligned} \quad (B13)$$

Thus, close to the weak coupling fixed point ($\Delta_K \approx 0$, $g_\gamma=1$), g_γ is marginal and Δ_K is relevant with dimension 1/2, while at the intermediate coupling fixed point (Δ_K finite and $g_\gamma \approx 0$), Δ_K is marginal and g_γ irrelevant. In addition, the flow equation for Δ_K in Eq(B13) determines the crossover scale, i.e. $\Delta_K(l^*) = O(1)$, where the renormalized values of the various amplitudes from the weak and intermediate coupling parts of the flows can be patched together.

Next, we will derive the RG flow equations close to the 2CK fixed point. This involves four usual steps. We will use a simple frequency shell RG, taking the unperturbed Hamiltonian to be H_0 with non-zero Δ_K and u ; note that as usual, certain coefficients will depend on the RG scheme. First the modes with magnitude of their frequency in the infinitesimal range between $\tau_c^{-1}(1-dl)$ and τ_c^{-1} are integrated out. Then the frequency cutoff is rescaled to its initial value τ_c^{-1} and the fields are rescaled by their scaling dimension, determined by their two point correlation functions, from which the RG eigenvalues of the various operators can be determined. Third, perturbative corrections to the flow equations are included from integrating out the high energy modes in the four fermion terms, for small $\bar{q}$, $\bar{y}$ and w . Thus, e.g. the contraction (or integration) of γ and β_I in the $\bar{y}$ -term contributes to the generation of Δ_+ . Finally, since at each step of this decimation process $i\gamma\beta_X$ will be generated, we have to rotate β_I and β_X so that it disappears. However, the corrections due to this rotation are negligible to the order at which we work. Following this

procedure we get the full set of RG equations describing the 2CK fixed point, and more generally for small Δ_+ , w , $\bar{y}$ and $\bar{q}$, but with general Δ_K ,

$$\begin{aligned}
\frac{d\Delta_+}{dl} &= \frac{1}{2}\Delta_+ + 2\bar{y}\Delta_K + \frac{wu}{\sqrt{2}(1+u^2)} + O(\bar{y}^2\Delta_+, w^2\Delta_+, wu\Delta_+, \Delta_+^3) \\
\frac{d\Delta_K}{dl} &= \frac{\Delta_K}{2} \left(1 - \frac{\Delta_K^2}{4\pi}\right) + 2\bar{y}\Delta_+ + O(\bar{y}^2\Delta_K, \Delta_+^3) \\
\frac{d\bar{q}}{dl} &= -\frac{\Delta_K^2}{8\pi}\bar{q} + O\left(\bar{q}^2\frac{\Delta_+}{\Delta_K}, \bar{y}^2, \frac{\Delta_+}{\Delta_K}\bar{q}\bar{y}\frac{\Delta_+}{\Delta_K}\right) \\
\frac{d\bar{y}}{dl} &= -\frac{\Delta_K^2}{8\pi}\bar{y} + O\left(\bar{q}^2\frac{\Delta_+}{\Delta_K}, \bar{y}^2\frac{\Delta_+}{\Delta_K}, \bar{q}\bar{y}\frac{\Delta_+}{\Delta_K}\right) \\
\frac{dw}{dl} &= -\frac{w}{2} + O(u\Delta_+) \\
\frac{du}{dl} &= -\frac{w\Delta_+}{4\pi}
\end{aligned} \tag{B14}$$

The constraint of Eq(B12) has been used; it has the effect of simplifying various messy terms that would appear in denominators from Eqs(B10) and (B11) evaluated at $|\omega| = 1/\tau_c$.²⁵ The above flow equations are identical to the weak coupling flow equations Eq(23), to lowest order in y , q , Δ_i and $q\Delta_i$ and $y\Delta_i$, with $i = +, -, 1$ or K . The higher order terms in the Δ_i 's are non-universal and depend on the RG procedure. In the flow equations above, we have selected a RG scheme via Eq(B12), such that the scaling dimensions of the various operators are given correctly to leading order in the both weak and intermediate coupling limits. The u dependence of the w term in $\frac{d\Delta_+}{dl}$ can, however, depend on the RG scheme, or, equivalently, the definition of u .

If w , Δ_+ and $\bar{y}$ are all small at some scale, then the $w\Delta_+$ and $\bar{y}\Delta_+$ terms can be ignored. The RG equations can then be integrated to yield the condition for lying on the critical manifold:

$$\Delta_+ + \frac{wu}{\sqrt{2}(1+u^2)} - \frac{2\bar{y}\Delta_K}{1 - \frac{\Delta_K^2}{4\pi}} \ln \left| \frac{\Delta_K^2}{4\pi} \right| = 0 \tag{B15}$$

to linear order in w , $\bar{y}$ and Δ_+ . In general, the parameters above should be evaluated in the regime of the crossover to intermediate coupling. [In the approximation of ignoring the neglected terms, the condition Eq(B15) can be used at any scale.] The role of w in the above equation is a typical example of the way an irrelevant operator can renormalize the value of a relevant one before decaying away. In this context, one should remember that the values of both Δ_K and Δ_+ should include corrections from the more strongly irrelevant operators that exist at higher energy scales but become negligibly small as we approach the fixed point, or as the flow moves away from weak coupling.

The effects of various symmetry breaking terms are discussed in Section IV.D. The symmetries of the important operators are summarized in Table II. If spin-flipping is allowed, it was shown in Section IV.D that the relevant processes involve joint electron-impurity hops which carry electronic spin. Their representation in terms of the bare fields involves ordering operators:

$$\begin{aligned}
\mathcal{H}_{sf} &= e^{i\pi(N_{ec}+N_{es})} \left\{ \Delta_{es} (\sigma_+ e^{i\Phi_c} - \sigma_- e^{-i\Phi_c}) (e^{i\Phi_{es}} - e^{-i\Phi_{es}}) \right. \\
&\quad - \Delta'_{es} i (\sigma_+ e^{i\Phi_c} - \sigma_- e^{-i\Phi_c}) (e^{i\Phi_{es}} + e^{-i\Phi_{es}}) \\
&\quad \left. + \Delta_s (\sigma_+ e^{i\Phi_c} - \sigma_- e^{-i\Phi_c}) (e^{i\Phi_s} - e^{-i\Phi_s}) \right\};
\end{aligned} \tag{B16}$$

but after refermionizing, these disappear with the choices made in Eq(B4), yielding, in the Majorana representation,

$$\mathcal{H}_{sf} = 2\sqrt{2\pi\tau_c}i (\Delta_{es}\delta\alpha_{es} + \Delta'_{es}\delta\beta_{es} + \Delta_s\delta\alpha_s). \tag{B17}$$

APPENDIX C: LEADING IRRELEVANT OPERATORS – COMPARISON TO CONFORMAL FIELD THEORY RESULTS

In this paper we have implemented an RG approach based on a bosonization method first developed by Emery and Kivelson¹⁸ which is inherently anisotropic in the channel pseudo-spin. We thus found a 2CK-like fixed point with an $O(2)$ symmetry, higher than the Z_2 interchange symmetry of the original impurity problem. Despite the fact that this

fixed point had lower symmetry than the $SU(2)$ symmetric one of the 2CK model solved by conformal field theory methods, we will show in this Appendix that *all* relevant, marginal and leading irrelevant operators expected from conformal field theory can be recovered from this model.¹²

To be specific, in Section IV.D the six dimension 1/2 relevant operators of the 2CK fixed point were analyzed (see Table II). Also, the only marginal, dimension 1, operator was found to be the conventional potential scattering term, $u\Psi_{ec}^\dagger\Psi_{ec}$, yielding $i\alpha_{ec}\beta_{ec}$.

In addition, there are four leading irrelevant, dimension 3/2, operators consistent with the symmetries of the problem we considered. The first, $\gamma\delta\alpha_c\beta_X$, corresponding to $\bar{q}$, does not break the $O(2)$ symmetry of the fixed point. This is essentially the leading irrelevant operator discussed in reference 10. In fact, there is no other operator with the same dimension,²⁶ invariant under $O(2)$ and the other discrete symmetries of the system, that does not involve even operators (Φ_{ec} , Φ_{es}), which naively decouple from the impurity. This can be checked directly from Table I, by taking all possible combinations of Majorana fermions in the channel pseudo-spin sector. Since the 2CK fixed point can be correctly described in terms of the free Majorana fermions we conclude that there is no other operator missing from our approach. This fact suggests that the operators that would distinguish the $O(2)$ from the $SU(2)$ symmetric fixed points have dimension higher than 3/2 and are thus more strongly irrelevant, in contrast with what Affleck *et al*¹¹ have claimed that a term breaking the $SU(2)$ down to an $O(2)$ symmetry has dimension 3/2.²⁷

The second leading irrelevant operator, $\delta\alpha_s\alpha_{es}\beta_{es}$, can be seen from Table I to be the only other operator with dimension 3/2 which respects the $O(2)$ as well as the discrete, symmetries of the system. Although it explicitly involves the Majorana fermions of the spin sector, it can be seen that this operator also satisfies the full spin $SU(2)$ symmetry by the fact that it is generated from a strictly spin $SU(2)$ invariant term, Δ_1 . Starting from Eq(5), one can first bosonize the fermion operators $c_{i\sigma}^\dagger$ directly into exponentials of combinations of Φ_{ec} , Φ_{es} , Φ_c and Φ_s . Then the Δ_1 operator becomes

$$\frac{\Delta_1}{4\pi\tau_c} \left\{ \sigma_+ \left[e^{-\frac{i}{2}(\Phi_{ec}+\Phi_{es}-\Phi_c-\Phi_s)} e^{\frac{i}{2}(\Phi_{ec}+\Phi_{es}+\Phi_c+\Phi_s)} \right. \right. \\ \left. \left. + e^{-\frac{i}{2}(\Phi_{ec}-\Phi_{es}-\Phi_c+\Phi_s)} e^{\frac{i}{2}(\Phi_{ec}-\Phi_{es}+\Phi_c-\Phi_s)} \right] + h.c. \right\} \quad (C1)$$

(multiplied by $\exp i\pi(N_{es} + N_{ec})$, the appropriate ordering operator discussed in Appendix B). Obviously, the leading operator of this term is the one appearing in Eq(13). However there are subleading corrections, coming from the operator product expansion of the various exponentials. One such term is

$$i \left(\sigma_+ e^{i\Phi_c} - \sigma_- e^{-i\Phi_c} \right) \frac{\partial \Phi_{es}}{\partial x} \sin \Phi_s, \quad (C2)$$

with a cutoff dependent coefficient proportional to Δ_1 . This is allowed by the full symmetries of the 2CK problem and will therefore be generated in the way outlined above or, equivalently, from the high energy scattering processes. At weak coupling, this term, with RG eigenvalue $-1/2 - 2q^2$, is always irrelevant. After performing the unitary transformation of Eq(28) and refermionizing, it becomes $\delta\alpha_s\alpha_{es}\beta_{es}$, with scaling dimension 3/2 close to the intermediate coupling fixed point. This operator contributes to the singular part of the impurity specific heat but not to the susceptibility of the conventional two channel, spin 1/2, Kondo model, thus leading to a *non-universal* Wilson ratio (The existence of two irrelevant operators at the 2CK fixed point was discussed by Affleck and Ludwig in Appendix D of reference 12). Other approaches to the 2CK problem^{10,18,28} assumed implicitly some higher symmetry²⁹ at the fixed point and therefore do not contain this second leading irrelevant operator (see reference 11 for more details).

The remaining two dimension 3/2 operators, $\gamma\delta\alpha_c\beta_I$ and $\delta\alpha_c\alpha_{ec}\beta_{ec}$, do *not* respect the $O(2)$ symmetry, but exist on the critical manifold with the Z_2 interchange symmetry, i.e. having set the relevant operator, corresponding to a linear combination of Δ_+ , w and $\bar{y}$, to zero — see Eq(B15). They were discussed in Section IV.C and correspond to couplings $\bar{y}$ and w . Interestingly, the first does not enter in the leading impurity specific heat correction of $O(T \ln T)$, since it couples to a dimension 2 irrelevant operator, g_γ , leading only to a $T^3 \ln T$ correction to the impurity specific heat.

More generally, in the case where Z_2 interchange symmetry is also broken, there are three more dimension 3/2 irrelevant operators, namely $\gamma\delta\alpha_{ec}\beta_{ec}$, $i\delta\frac{\partial\beta_s}{\partial x}$ and $i\delta\frac{\partial\beta_c}{\partial x}$. However, in this case *two* relevant operators must be tuned to be on the critical manifold.

Finally, for completeness, we summarize the leading irrelevant operators for the additional non-Fermi liquid fixed point discussed in Section V. Apart from $\bar{q}$ and $\bar{y}$, the operators $\gamma\beta_I\alpha_{ec}\beta_{ec}$ and $\gamma\beta_X\alpha_{ec}\beta_{ec}$ also have scaling dimension 3/2. Again, only the last two contribute to $O(T \ln T)$ to the impurity specific heat. As in the above case, in the absence of Z_2 symmetry, three additional dimension 3/2 operators will be present, namely $i\gamma\frac{\partial\alpha_c}{\partial x}$, $\gamma\delta\alpha_{ec}\beta_{ec}$ and $\gamma\alpha_c\beta_c\beta_s$.

Fermion	Site Interchange (Z_2)	Time Reversal	Spin Reversal	Spin Conservation mod 2
γ	+	+	+	+
δ	-	-	+	+
α_c	-	+	+	+
β_c	+	-	+	+
α_s	-	+	-	+
β_s	+	-	+	+
α_{ec}	+	+	+	-
β_{ec}	+	-	+	-
α_{es}	+	+	-	-
β_{es}	+	-	+	-

TABLE I. Transformation properties of Majorana fermions with respect to the discrete symmetries of the system. Electron number conservation implies that the Bose field Φ_{ec} can be shifted by an arbitrary amount, which means that only the combination $\alpha_{ec}\beta_{ec}$ can appear. In the presence of conservation of the z -component of the spin, Φ_{es} can also be shifted by an arbitrary constant. If z -spin were only conserved mod 2, then just a discrete symmetry would remain; this is listed in the last column of the table as a simple way to distinguish the different symmetries of the e operators.

Relevant Operators	Site Interchange (Z_2)	Time Reversal	Spin Reversal	Spin Conservation mod 2
$i\delta\alpha_c$	+	+	+	+
$i\delta\gamma$	-	+	+	+
$i\delta\beta_X$	-	-	+	+
$i\delta\alpha_s$	+	+	-	+
$i\delta\alpha_{es}$	-	+	-	-
$i\delta\beta_{es}$	-	-	+	-

TABLE II. The six independent relevant operators at 2CK fixed point and their transformation properties (for last column, see note in Table I). Note that $i\delta\beta_s$ and $i\delta\beta_c$ are not listed here but rather the combination $i\delta\beta_X$ which has the same symmetries as the other two. (See discussion in Section IV.D).

-
- ¹ P. Nozières and A. Blandin, J. Phys. (Paris) **41**, 193 (1980).
² D. L. Cox, Phys. Rev. Lett. **59**, 1240 (1987).
³ K. Vladár and A. Zawadowski, Phys. Rev. B **28**, 1564 (1983); B **28**, 1582 (1983); B **28**, 1596 (1983).
⁴ D. C. Ralph and R. A. Buhrman, Phys. Rev. Lett. **69**, 2118 (1992).
⁵ D. C. Ralph, A. W. W. Ludwig, Jan von Delft and R. A. Buhrman, Phys. Rev. Lett. **72**, 1064 (1994).
⁶ I. Affleck and A. W. W. Ludwig, Phys. Rev. Lett. **67**, 3160 (1991); A. W. W. Ludwig and I. Affleck, Phys. Rev. B **48**, 7297 (1993).
⁷ N. S. Wingreen, B. L. Altshuler and Y. Meir, Phys. Rev. Lett. **75**, 769 (1995); D. C. Ralph, A. W. W. Ludwig, Jan von Delft and R. A. Buhrman, Phys. Rev. Lett. **75**, 770 (1995).
⁸ K. Vladár, G. T. Zimányi and A. Zawadowski, Phys. Rev. Lett. **56**, 286 (1985).
⁹ A. Murumatsu and F. Guinea, Phys. Rev. Lett. **57**, 2337 (1986).

- ¹⁰ A. M. Sengupta and A. Georges, Phys. Rev. B **49**, 10020 (1994).
¹¹ I. Affleck, A. W. W. Ludwig, H. B. Pang and D. L. Cox, Phys. Rev. B **45**, 7918 (1992).
¹² I. Affleck and A. W. W. Ludwig, Nucl. Phys. B **360**, 641 (1991).
¹³ K. Yamada, A. Sakurai and M. Takeshige, Prog. Theor. Phys. **70**, 73 (1970).
¹⁴ A. L. Moustakas and D. S. Fisher, Phys. Rev. B **51**, 6908 (1995).
¹⁵ It turns out that the effects of this artificial symmetry remain even if it holds modulo 2 (i.e. $N_1 - N_2 \rightarrow N_1 - N_2 \pm 2$).
¹⁶ In the case of intersite separation that is much larger than the inverse Fermi wavevector, both y and the momentum dependence of scattering away from the Fermi surface can be regarded as small¹⁴.
¹⁷ Yu. Kagan and N. V. Prokof'ev, Sov. Phys. JETP **69**, 836 (1989) (Zh. Eksp. Teor. Fiz. **96**, 1473 (1989)).
¹⁸ V. J. Emery and S. Kivelson, Phys. Rev. B **46**, 10812 (1992).
¹⁹ Throughout this paper, we will be careful to keep all terms whose presence, or generation under renormalization, plays an important role either for the crossover when some of the

hopping terms are small, or for the behavior near the 2CK fixed point. The other terms, which we neglect, should only give rise to no more than (generally non-universal) order one pre-factors.

²⁰ G. Toulouse, Phys. Rev. B **2**, 270 (1970).

²¹ P. Nozières, *private communication*.

²² M. Fabrizio, A. O. Gogolin and P. Nozières, Phys. Rev. Lett. **74**, 4503 (1995).

²³ P. Coleman and A. J. Schofield, preprint (cond-mat 9504084).

²⁴ A similar situation of an apparent extra, but redundant relevant operator, appears in conventional ϕ^4 theory: the ϕ^3 operator, in the presence of a magnetic field term $h\phi$ can be transformed away by a non-singular (albeit non-linear) transformation of the field ϕ .

²⁵ It should be noted, as pointed out in the main text that, due to the inherent anisotropy of the RG scheme employed here, the fixed point Hamiltonian will *not* have the channel $SU(2)$ symmetry of the 2CK problem.

²⁶ We are grateful to Anirvan Sengupta for confirming this point to us. More details can be found in A. Georges and A. M. Sengupta, Phys. Rev. Lett. **74**, 2808 (1995).

²⁷ However, in the four channel Kondo model, there are *two* distinct leading irrelevant operators consistent with $O(2)$ symmetry. See M. Fabrizio and A. O. Gogolin, Phys. Rev. B **50**, 17732 (1994).

²⁸ A. M. Tsvelick and P. B. Wiegmann, J. Stat. Phys. **38**, 125 (1985).

²⁹ A. M. Sengupta, *private communication*.